

Technical Data Sheet


Purified Mouse Anti-Human CD196 (CCR6)**Product Information**

Material Number:	559560
Alternate Name:	CCR6
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	11A9
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The monoclonal antibody 11A9 reacts with the human CC chemokine receptor, CCR6. CCR6 (previously known as BN-1, CKR-L3, DRY6, GPR-CY4 and STRL22), a seven-transmembrane, G-protein-coupled receptor, is the specific receptor for CC chemokine MIP-3a/LARC/Exodus. It has been shown that CCR6 mRNA is expressed mainly in lymphoid tissues including spleen, lymph nodes, thymus, appendix. CCR6 mRNA was also detected in peripheral T- and B-lymphocytes and in CD34-derived dendritic cells. The human CCR6 gene, unlike other CCR genes, has been mapped to chromosome 6q27. The immunogen used to generate 11A9 hybridoma was KLH-conjugated N-terminus peptides of human CCR6. It does not cross-react with human CCR1, CCR2, CCR3, CCR4, CCR5, CCR7, CCR8, CCR9, CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 transfectants. This antibody is NOT a neutralizing antibody. CCR6 had been clustered as CD196 in the VIIIth HLDA workshop.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Expression of human CCR6 on peripheral lymphocytes detected by purified 11A9.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4° C.

Application Notes**Application**

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

The purified 11A9 antibody (Cat. No. 559560) can be used for the immunofluorescent staining and flow cytometric analyses of human leukocytes (see image) and cell lines that express CCR6.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



BD

BD Biosciences

A multistep staining procedure is recommended to amplify immunofluorescent signals for the flow cytometric analysis of human CCR6 expression:

Step 1: Incubate the cells with 0.1 - 1 µg of purified 11A9 antibody at 4°C for 15-20 minutes. Wash cells two times with staining medium containing sodium azide (e.g., Dulbecco's PBS or tissue culture medium [without phenol red and biotin] with 0.09% sodium azide and 2% heat-inactivated FCS or 0.2% BSA).

Step 2: Incubate the cells with 0.25 µg of biotinylated goat anti-mouse Ig (Cat. No. 553999) at 4°C for 20 minutes. Wash cells two times.

Step 3: Incubate the cells with ≤ 0.06 mg of streptavidin-phycoerythrin (Cat. No. 554061) at 4°C for 20 minutes. Wash two times. Resuspend cells in staining medium and analyze stained cells with a BD FACScan™ Flow Cytometer (BD Biosciences, San Jose, CA) using appropriate specificity and compensation controls.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554061	Streptavidin PE	0.5 mg	(none)
555746	Purified Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21
553999	Biotin Goat Anti-Mouse Ig	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

- Baba M, Imai T, Nishimura M, et al. Identification of CCR6, the specific receptor for a novel lymphocyte-directed CC chemokine LARC. *J Biol Chem.* 1997; 272(23):14893-14898.(Biology)
- Greaves DR, Wang W, Dairaghi DJ, et al. CCR6, a CC chemokine receptor that interacts with macrophage inflammatory protein 3alpha and is highly expressed in human dendritic cells. *J Exp Med.* 1997; 186(6):837-844.(Biology)
- Liao F, Alderson R, Su J, Ullrich SJ, Kreider BL, Farber JM. STRL22 is a receptor for the CC chemokine MIP-3alpha. *Biochem Biophys Res Commun.* 1997; 236(1):212-217.(Biology)
- Liao F, Lee HH, Farber JM. Cloning of STRL22, a new human gene encoding a G-protein-coupled receptor related to chemokine receptors and located on chromosome 6q27. *Genomics.* 1997; 40(1):175-180.(Biology)
- Liao F, Rabin RL, Smith CS, Sharma G, Nutman TB, Farber JM. CC-chemokine receptor 6 is expressed on diverse memory subsets of T cells and determines responsiveness to macrophage inflammatory protein 3 alpha. *J Immunol.* 1999; 162(1):186-194.(Biology)
- Power CA, Church DJ, Meyer A, et al. Cloning and characterization of a specific receptor for the novel CC chemokine MIP-3alpha from lung dendritic cells. *J Exp Med.* 1997; 186(6):825-835.(Biology)
- Zaballos A, Varona R, Gutierrez J, Lind P, Marquez G. Molecular cloning and RNA expression of two new human chemokine receptor-like genes. *Biochem Biophys Res Commun.* 1996; 227(3):846-853.(Biology)