

Technical Data Sheet

Fixable Viability Stain 700

Product Information


Material Number: 564997

Size: 100 µg

Description

BD Horizon™ Fixable Viability Stain 700 (FVS700) is useful for discrimination of viable from non-viable mammalian cells in multicolor flow cytometric applications. This dye reacts with and covalently binds to cell-surface and intracellular amines. Permeable plasma cell membranes, such as those present in necrotic cells, allow for the intracellular diffusion and covalent binding of the dye to the higher levels of total cellular amines when compared to living cells with impermeable plasma membranes. Therefore, necrotic cells present in a typical *in vitro* assay label with higher levels of dye increasing their FVS700 fluorescence intensities 10-20 fold over that of viable cells. The FVS700-labeled cells can be fixed with formaldehyde for downstream decontamination, freezing and/or permeabilization and subsequent intracellular staining while maintaining stable viability stain fluorescence.

BD Horizon™ Fixable Viability Stain 700 is excited by the Red laser (with an excitation maximum of 657 nm), and has a fluorescence emission maximum of 700 nm.



Flow cytometric analysis of human Jurkat cells stained with BD Horizon™ Fixable Viability Stain 700. Cells from the human Jurkat (Acute T cell leukemia, ATCC TIB-152) cell line were treated with 0.025% DMSO (Left Panel) or 5 µM camptothecin (Right Panel) for 16 hours and then stained with BD Horizon™ Fixable Viability Stain 700 (Cat. No. 564997) in serum-free buffer. The cells were then either left unfixed (solid line histograms) or fixed in BD Cytotfix™ Fixation Buffer (Cat. No. 554655) and permeabilized in BD Phosflow™ Perm/Wash Buffer I (Cat. No. 557885, dashed line histograms). Histograms were derived from gated events with the forward and side light-scattering characteristics of Jurkat cells. Flow cytometric analysis was performed using a BD LSRFortessa™ Flow Cytometry System. Please note that FVS700 is also compatible with BD Phosflow™ Perm Buffer III (Cat. No. 558050) or BD Pharmingen™ Transcription Factor Buffer Set (Cat. No. 562574/562725). Fixable Viability Stain 700 has been tested on mouse (data not shown).

Application Notes

Application

Flow cytometry	Tested During Development
Intracellular staining (flow cytometry)	Tested During Development

Recommended Assay Procedure:

Preparation

Bring FVS700 dye powder and 310 µl of fresh cell culture-grade Dimethyl Sulfoxide (DMSO; eg, Sigma D2650) to room temperature. Add 310 µl of DMSO and vortex solution well. Inspect the solution and repeat vortex until the stock dye has fully dissolved. This is the Stock Solution.

Storage

Upon arrival, store the dry dye desiccated and protected from light at -80°C until use. After reconstitution with DMSO, store the Stock Solution at -20°C in small aliquots. Do not use reconstituted dye after 90 days of storage. Please discard the dye solution after 90 days post reconstitution with DMSO.

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean
877.232.8995 866.979.9408 32.2.400.98.95 0120.8555.90 65.6861.0633 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

564997 Rev. 2



Cytometry Requirements

Red laser-equipped Flow Cytometers (eg, BD FACSCanto™ II, BD LSRFortessa™, or BD LSR™ II) can be used. This dye can be read out of filters commonly used for BD Horizon™ APC-R700 or Alexa Fluor® 700 (eg, 730/45- or 712/21-nm filter). Fluorescence compensation is best achieved using stained and unstained samples of the cells of interest.

Procedure

Fixable Viability Stain 700 labeling of cells

1. Prepare cells for flow cytometric staining using sodium azide-free buffers.
2. Wash cells one time in sodium azide- and protein-free Dulbecco's Phosphate Buffered Saline (1X DPBS).
3. Resuspend cells at $1-10 \times 10^6$ cells/ml in sodium azide- and protein-free 1X DPBS.
4. Add 1 μ l of BD Horizon™ Fixable Viability Stain 700 Stock Solution for each 1 ml of cell suspension (1:1000) and vortex immediately.
 - a. *Note:* We recommend titrating the dye for optimal performance as different cell types and different applications can result in a wide degree of variability in staining. Please see Note 1 below for guidance on recommended ranges.
5. Incubate the mixture for 10-15 minutes at room temperature or 2-8°C protected from light.
 - a. *Optional:* Alternatively, incubate mixtures at 37°C for 5-7 minutes.
6. Wash cells twice with 2 ml of BD Pharmingen™ Stain Buffer (FBS) (Cat. No. 554656) or the equivalent.
7. Decant the supernatant and gently mix to disrupt the cell pellet.
8. Resuspend the cells in Stain Buffer (FBS) or equivalent.
9. Stain, fix and permeabilize cells as desired for downstream applications.

Notes:

1. Each user should determine the optimal concentrations of reagents, cells, and conditions for the assay of interest. We recommend titrating the reagent in early experiments to obtain optimal results. The following are ranges that we have found to work for various cell systems:
 - a. Erythrocyte-Lysed Whole Blood Cells: 1:1,000 from the Stock Solution.
 - b. Primary Cells: 1:1,000 - 1:4,000 from the Stock Solution.
 - c. Cell Lines: 1:4,000 - 1:10,000 from the Stock Solution.
2. The reactivity of the free dye is quenched by washing with buffer containing protein (eg, FBS or BSA).
3. Cells may be stained in bulk prior to freezing or staining with fluorescent antibodies.
4. BD Horizon™ Fixable Viability Stain 700 can be used in intracellular staining assays that require fixation with formaldehyde and permeabilization with methanol and detergents such as those used for BD Phosflow™ staining (eg, BD Phosflow™ Perm Buffer III, Cat. No. 558050), intracellular cytokine staining (eg, BD Cytotfix/Cytoperm™ Fixation/Permeabilization Solution Kit, Cat. No. 554714), or transcription factor staining (eg, BD Pharmingen™ Transcription Factor Buffer Set, Cat. No. 562574/562725).
5. Apoptotic cells can show variable staining. We recommend co-staining with other fluorescent probes such as Annexin V FITC (Cat. No. 556419) if further analysis is desired for the apoptotic cells.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 mL	(none)
554655	Fixation Buffer	100 mL	(none)
557885	Perm/Wash Buffer I	125 mL	(none)
558050	Perm Buffer III	125 mL	(none)
562574	Transcription Factor Buffer Set	100 Tests	(none)
562725	Transcription Factor Buffer Set	25 Tests	(none)
554714	BD Cytotfix/Cytoperm™ Fixation/Permeabilization Kit	250 Tests	(none)
556419	FITC Annexin V	200 Tests	(none)
554657	Stain Buffer (BSA)	500 mL	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
3. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
4. Before staining with this reagent, please confirm that your flow cytometer is capable of exciting the fluorochrome and discriminating the resulting fluorescence.
5. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Abrams B, Diwu Z, Guryev O, et al. 3-Carboxy-6-chloro-7-hydroxycoumarin: a highly fluorescent, water-soluble violet-excitable dye for cell analysis. *Anal Biochem.* 2009; 386(2):262-269. (Methodology)

Burmeister Y, Lischke T, Dahler AC, et al. ICOS controls the pool size of effector-memory and regulatory T cells. *J Immunol.* 2008; 180(2):774-782. (Methodology)

Charles ED, Green RM, Marukian S, et al. Clonal expansion of immunoglobulin M+CD27+ B cells in HCV-associated mixed cryoglobulinemia. *Blood.* 2008; 111(3):1344-1356. (Methodology)

Perfetto SP, Chattopadhyay PK, Lamoreaux L, et al. Amine-reactive dyes for dead cell discrimination in fixed samples. *Curr Protoc Cytom.* 2010; 9(9.34) (Methodology)

Perfetto SP, Chattopadhyay PK, Lamoreaux L, et al. Amine reactive dyes: an effective tool to discriminate live and dead cells in polychromatic flow cytometry. *J Immunol Methods.* 2006; 313(1-2):199-208. (Methodology)