

## Technical Data Sheet

**PerCP-Cy™ 5.5 Rat IgG2a, κ Isotype Control****Product Information**

|                         |                                                           |
|-------------------------|-----------------------------------------------------------|
| <b>Material Number:</b> | <b>550765</b>                                             |
| <b>Size:</b>            | 0.1 mg                                                    |
| <b>Concentration:</b>   | 0.2 mg/ml                                                 |
| <b>Clone:</b>           | R35-95                                                    |
| <b>Immunogen:</b>       | Mouse Pooled Immunoglobulin                               |
| <b>Isotype:</b>         | Rat (LOU) IgG2a, κ                                        |
| <b>Storage Buffer:</b>  | Aqueous buffered solution containing ≤0.09% sodium azide. |

**Description**

The R35-95 hybridoma was generated by hybridization of Y3 myeloma cells with spleen cells from LOU rats immunized with mouse immunoglobulins. The R35-95 hybridoma produces rat IgG2a, κ immunoglobulin that has no measurable reactivity with mouse immunoglobulins. The R35-95 immunoglobulin was selected as an isotype control following screening for low background binding on a variety of mouse and human tissues.

**Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with PerCP-Cy5.5 under optimum conditions, and unconjugated antibody and free PerCP-Cy5.5 were removed. Storage of PerCP-Cy5.5 conjugates in unoptimized diluent is not recommended and may result in loss of signal intensity. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

**Application Notes****Application**

|                 |                  |
|-----------------|------------------|
| Flow cytometry  | Routinely Tested |
| Isotype control | Routinely Tested |

**Recommended Assay Procedure:**

An isotype control should be used at the same concentration as the antibody of interest (e.g., ≤ 1 µg/million cells).

PerCP has been reported to undergo significant photobleaching, the magnitude of which increases as laser power is increased or beam focus is narrowed. For tandem conjugates incorporating PerCP (e.g., PerCP-Cy5.5), the excitation and emission properties of PerCP and the kinetics of energy exchange between the fluorochromes of the tandem dye may limit their effectiveness on high-speed and/or sorting flow cytometers. Therefore, for third-color flow cytometric analysis using ≥ 25-mW laser power, we recommend PE-Cy5 (formerly BD Cy-Chrome™)-conjugated reagents.

It is recommended that a 712/20-nm band-pass filter be used with stream-in-air instruments such as the BD FACStar™ and BD FACSVantage™ Flow Cytometry Systems.

**Product Notices**

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Cy is a trademark of Amersham Biosciences Limited. This conjugated product is sold under license to the following patents: US Patent Nos. 5,486,616; 5,569,587; 5,569,766; 5,627,027.
4. PerCP-Cy5.5-labelled antibodies can be used with FITC- and R-PE-labelled reagents in single-laser flow cytometers with no significant spectral overlap of PerCP-Cy5.5, FITC, and R-PE fluorescence.
5. PerCP-Cy5.5 is optimized for use with a single argon ion laser emitting 488-nm light. Because of the broad absorption spectrum of the tandem fluorochrome, extra care must be taken when using dual-laser cytometers, which may directly excite both PerCP and Cy5.5™. We recommend the use of cross-beam compensation during data acquisition or software compensation during data analysis.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
7. This product is subject to proprietary rights of Amersham Biosciences Corp. and Carnegie Mellon University and made and sold under license from Amersham Biosciences Corp. This product is licensed for sale only for research. It is not licensed for any other use. If you require a commercial license to use this product and do not have one return this material, unopened to BD Biosciences, 10975 Torreyana Rd, San Diego, CA 92121 and any money paid for the material will be refunded.
8. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

**BD Biosciences**

[bdbiosciences.com](http://bdbiosciences.com)

|                      |               |               |              |                     |                                |
|----------------------|---------------|---------------|--------------|---------------------|--------------------------------|
| <b>United States</b> | <b>Canada</b> | <b>Europe</b> | <b>Japan</b> | <b>Asia Pacific</b> | <b>Latin America/Caribbean</b> |
| 877.232.8995         | 888.268.5430  | 32.53.720.550 | 0120.8555.90 | 65.6861.0633        | 0800.771.7157                  |

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## References

Greimers R, Trebak M, Moutschen M, Jacobs N, Boniver J. Improved four-color flow cytometry method using fluo-3 and triple immunofluorescence for analysis of intracellular calcium ion ( $[Ca^{2+}]_i$ ) fluxes among mouse lymph node B- and T-lymphocyte subsets. *Cytometry*. 1996; 23(3):205-217. (Biology)

Shapiro HM. *Practical Flow Cytometry, 3rd Edition*. New York: Wiley-Liss, Inc; 1995:280-281. (Biology)