

Technical Data Sheet

Oligo Mouse Anti-Human Chromogranin A

Product Information

Material Number:	940512
Size:	25 Tests
Clone:	S21-537
Alternative Name:	CGA; CgA; CHGA; Chromogranin-A; CMGA; SP-I
Reactivity:	Tested in Development: Human
Isotype:	Mouse IgG1, κ
Application:	Single Cell 3' Sequencing Intracellular CITE-seq (Tested During Development)
Barcode Sequence:	GAAAGCAGGTTGGCCGTGTTAAGCGTGATCGAAGT
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.
Regulatory Status:	RUO

Description

Chromogranin A (CGA) is a member of the granin family of regulated secretory proteins that are found in secretory granules in endocrine and neuroendocrine cells and released in response to extracellular stimulation. Intracellularly, granins are important for targeting peptide hormones and neurotransmitters by their ability to aggregate in the low pH, high calcium environment of the trans-Golgi network. Extracellularly, peptides formed from proteolytic processing of granins regulate hormone secretion. CGA is a prohormone that can be cleaved into several biologically active peptides, such as pancreastatin, β -granin, vasostatin, catestatin, and parastatin. β -granin is an N-terminal fragment of CGA, while pancreastatin and catestatin are processed from the central region of CGA. Cells of the adrenal medulla, anterior pituitary, cerebral cortex as well as beta cells of the pancreas and a variety of tumor cell lines express CGA. The expression of CGA can be used to monitor the pancreatic differentiation of pluripotent stem cells.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography and conjugated to BD® AbSeq oligonucleotide under optimal conditions.

Recommended Assay Procedure

Put all BD® AbSeq reagents to be pooled into a Latch Rack for 500 μ L Tubes (Thermo Fisher Scientific Cat. No. 4900). Arrange the tubes so that they can be easily uncapped and re-capped with an 8-Channel Screw Cap Tube Capper (Thermo Fisher Scientific Cat. No. 4105MAT) and the reagents aliquoted with a multi-channel pipette. BD® AbSeq tubes should be centrifuged for = 30 seconds at 400 $\times$ g to ensure removal of any content in the cap/tube threads prior to the first opening.

When using BD® AbSeq intracellular markers with the Single Cell 3' Sequencing Intracellular CITE-seq, cells must first be fixed and permeabilized using the BD Rhapsody™ Intracellular AbSeq Buffer Kit before the antibody-oligo can bind to the protein. Refer to the list of required companion products below and see BD Rhapsody™ System Single-Cell Labelling with BD® AbSeq Ab-Oligos for Intracellular CITE-seq (Doc ID: 23-24464) for the complete BD® AbSeq intracellular multiomics staining protocol. Contact your local Field Application Specialist (FAS) for additional guidance.

Use standard laboratory safety protocols. Read and understand the safety data sheets (SDSs) before handling chemicals. To obtain SDSs, go to regdocs.bd.com or contact BD Biosciences technical support at scomix@bdscomix.bd.com.

Warning: All biological specimens and materials contacting them are considered biohazardous. Handle as if capable of transmitting infection and dispose of with proper precautions in accordance with federal, state, and local regulations. Never pipette by mouth. Wear suitable protective clothing, eyewear, and gloves.

Suggested Companion Products

Catalog Number	Name	Size
570742	Intracellular AbSeq Buffer Kit	1 Each
570750	AbSeq Enhancer Kit	1 Each
570908	OMICS-Guard Sample Preservation Buffer Kit	12 Each

570751	RNase Inhibitor	1 Each
633801	Whole Transcriptome Analysis (WTA) Amplification Kit	1 Each
554656	Stain Buffer (FBS)	500 mL
564219	Human BD Fc Block™	50 µg
664887	Enhanced Cartridge Reagent Kit	1 Each
633733	Cartridge Kit	1 Each
666262	8-Lane Cartridge	1 Each
625970	Immune Discovery Panel	5 Tests
633773	cDNA Kit	1 Each
633781	Hu Single Cell Sample Multiplexing Kit	1 Each
570911	OMICS-Guard Sample Preservation Buffer	50 mL
633707	Express Single-Cell Analysis System Package	1 EA
633701	Single-Cell Analysis System	1 EA

Product Notices

1. Please refer to www.bdbiosciences.com/us/s/resources for technical protocols.
2. This reagent has been pre-diluted for use at the recommended volume per test. Typical use is 2 µl for 1 × 10⁶ cells in a 200-µl staining reaction.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
6. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
7. Please refer to bd.com/genomics-resources for technical protocols.
8. Illumina is a trademark of Illumina, Inc.
9. For U.S. patents that may apply, see bd.com/patents.

References

Hendy GN, Bevan S, Mattei MG, Mouland AJ. Chromogranin A. Clin Invest Med. 1995; 18(1):47-65. (Biology).

Taylor CV, Taupenot L, Mahata SK. Formation of the catecholamine release-inhibitory peptide catestatin from chromogranin A. Determination of proteolytic cleavage sites in hormone storage granules. Clin Invest Med. 2000; 275(30):22905-22915. (Biology).

Kroon E, Martinson LA, Kadoya K, Bang AG, et al. Pancreatic endoderm derived from human embryonic stem cells generates glucose-responsive insulin-secreting cells in vivo. Nat Biotechnol. 2008; 26(4):443-452. (Biology).

D'Amour KA, Bang AG, Eliazar S, et al. Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells. Nat Biotechnol. 2006; 24(12):1481-1483. (Biology).

Mouland AJ, Bevan S, White JH, Hendy GN. Human chromogranin A gene. Molecular cloning, structural analysis, and neuroendocrine cell-specific expression. J Biol Chem. 1994; 269(9):6918-6926. (Biology).

Schulz TC, Young HY, Agulnick AD et al. A scalable system for production of functional pancreatic progenitors from human embryonic stem cells. PLoS ONE. 7(5)(Biology).

Loh YP, Cheng Y, Mahata SK, Corti A, Tota B. Chromogranin A and derived peptides in health and disease. J Mol Neurosci. 2012; 48(2):347-356. (Biology).

Mendieta I, Nuñez-Anita RE, Pérez-Sánchez G, et al. Effect of A549 neuroendocrine differentiation on cytotoxic immune response.. Endocr Connect. 2018; 7(5):791-802. (Clone-specific: Intracellular Staining/Flow Cytometry).

BD Biosciences

bdbiosciences.com

United States 877.232.8995	Canada 888.268.5430	Europe 32.53.720.550	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 0800.771.7157
-------------------------------	------------------------	-------------------------	-----------------------	------------------------------	--

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other

©2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.