

Technical Data Sheet

Oligo Mouse Anti-T-bet

Product Information

Material Number:	940515
Size:	25 Tests
Clone:	O4-46
Alternative Name:	T-box expressed in T cells; TBX21; T-box 21; TBLYM
Reactivity:	Tested in Development: Human, Mouse
Isotype:	Mouse IgG1, κ
Application:	Single Cell 3' Sequencing Intracellular CITE-seq (Tested During Development)
Barcode Sequence:	TGTAAAGATGTCGGGCTATGGGTCAGTATGTATTGG
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.
Regulatory Status:	RUO

Description

The O4-46 monoclonal antibody specifically binds to human and mouse T-bet. T-bet (T-box gene expressed in T cells) is a master regulatory transcription factor that is also known as TBX21 (T-box21) and TBLYM (T-box transcription factor, expressed in lymphocytes). Human (535 amino acids; 58.3 kDa predicted molecular mass) and mouse (530 amino acids; 57.7 kDa) T-bet proteins are encoded by the human TBX21 (chromosome 17) and mouse Tbx21 (chromosome 11) genes. The human and mouse T-bet protein amino acid sequences are 88% homologous. Human and mouse T-bet proteins share a highly conserved (98% homologous amino acid sequences) T-box protein domain that is centrally located and mediates binding to DNA. T-bet is expressed by and activates transcriptional activities within hematopoietic cells including stem cells, NK and NKT cells and subsets of thymocytes, primed/activated CD4⁺ T cells, CD8⁺ T cells and $\gamma\delta$ T cells, B cells, and dendritic cells. Interferon-gamma (IFN- γ), interleukin-27 (IL-27), and IL-12 act on peripheral antigen-triggered (TCR-signaling) T cells to increase T-bet expression. With respect to T helper lymphocytes, T-bet directs the differentiation of naïve CD4⁺ precursor T cells to become Th1-like effector and memory cells. T-bet accomplishes this by activating Th1 genetic programs (including epigenetic modifications) while repressing opposing T helper subset programs. T-bet controls the upregulated expression of the Th1 signature cytokine, IFN- γ , the IL-12R β 2 subunit and the Runx3 transcription factor and can repress the function of other transcriptional regulators, such as GATA-3 (master regulator of Th2 development) and the expression of other cytokines including IL-2, IL-4 and IL-5.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography and conjugated to BD® AbSeq oligonucleotide under optimal conditions.

Recommended Assay Procedure

Put all BD® AbSeq reagents to be pooled into a Latch Rack for 500 μ L Tubes (Thermo Fisher Scientific Cat. No. 4900). Arrange the tubes so that they can be easily uncapped and re-capped with an 8-Channel Screw Cap Tube Capper (Thermo Fisher Scientific Cat. No. 4105MAT) and the reagents aliquoted with a multi-channel pipette. BD® AbSeq tubes should be centrifuged for = 30 seconds at 400 \times g to ensure removal of any content in the cap/tube threads prior to the first opening.

When using BD® AbSeq intracellular markers with the Single Cell 3' Sequencing Intracellular CITE-seq, cells must first be fixed and permeabilized using the BD Rhapsody™ Intracellular AbSeq Buffer Kit before the antibody-oligo can bind to the protein. Refer to the list of required companion products below and see BD Rhapsody™ System Single-Cell Labelling with BD® AbSeq Ab-Oligos for Intracellular CITE-seq (Doc ID: 23-24464) for the complete BD® AbSeq intracellular multiomics staining protocol. Contact your local Field Application Specialist (FAS) for additional guidance.

Use standard laboratory safety protocols. Read and understand the safety data sheets (SDSs) before handling chemicals. To obtain SDSs, go to regdocs.bd.com or contact BD Biosciences technical support at scomix@bdscomix.bd.com.

Warning: All biological specimens and materials contacting them are considered biohazardous. Handle as if capable of transmitting infection and dispose of with proper precautions in accordance with federal, state, and local regulations. Never pipette by mouth. Wear suitable protective clothing, eyewear, and gloves.

Suggested Companion Products

Catalog Number	Name	Size
570742	Intracellular AbSeq Buffer Kit	1 Each
570750	AbSeq Enhancer Kit	1 Each
570908	OMICS-Guard Sample Preservation Buffer Kit	12 Each
570751	RNase Inhibitor	1 Each
633801	Whole Transcriptome Analysis (WTA) Amplification Kit	1 Each
554656	Stain Buffer (FBS)	500 mL
564219	Human BD Fc Block™	50 µg
664887	Enhanced Cartridge Reagent Kit	1 Each
633733	Cartridge Kit	1 Each
666262	8-Lane Cartridge	1 Each
625970	Immune Discovery Panel	5 Tests
633773	cDNA Kit	1 Each
633781	Hu Single Cell Sample Multiplexing Kit	1 Each
570911	OMICS-Guard Sample Preservation Buffer	50 mL
633707	Express Single-Cell Analysis System Package	1 EA
633701	Single-Cell Analysis System	1 EA

Product Notices

1. Please refer to www.bdbiosciences.com/us/s/resources for technical protocols.
2. This reagent has been pre-diluted for use at the recommended volume per test. Typical use is 2 µl for 1 × 10⁶ cells in a 200-µl staining reaction.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
6. Species cross-reactivity detected in product development may not have been confirmed on every format and/or application.
7. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
8. Please refer to bd.com/genomics-resources for technical protocols.
9. Illumina is a trademark of Illumina, Inc.
10. For U.S. patents that may apply, see bd.com/patents.

References

Reis BS, Rogoz A, Costa-Pinto FA, Taniuchi I, Mucida D. Mutual expression of the transcription factors Runx3 and ThPOK regulates intestinal CD4(+) T cell immunity. *Nat Immunol.* 2013; 14(3):271-280. (Clone-specific: Flow cytometry).

Trotta E, Bessette PH, Silveria SL, et al. A human anti-IL-2 antibody that potentiates regulatory T cells by a structure-based mechanism. *Nat Med.* 2018; 24(7):1005-1014. (Clone-specific: Flow cytometry).

Li S, Sullivan NL, Rouphael N, et al. Metabolic Phenotypes of Response to Vaccination in Humans. *Cell.* 2017; 169(5):862-877.e17. (Clone-specific: Flow cytometry).

Ponzetta A, Carriero R, Carnevale S, et al. Neutrophils Driving Unconventional T Cells Mediate Resistance against Murine Sarcomas and Selected Human Tumors. *Cell.* 2019; 178(2):346-360.e24. (Clone-specific: Flow cytometry).

Cella M, Gamini R, Sécca C, et al. Subsets of ILC3-ILC1-like cells generate a diversity spectrum of innate lymphoid cells in human mucosal tissues. *Nat Immunol.* 2019; 20(8):980-991. (Clone-specific: Flow cytometry).

Kawabe T, Yi J, Kawajiri A, et al. Requirements for the differentiation of innate T-bethigh memory-phenotype CD4+ T lymphocytes under steady state. *Nat Commun.* 2020; 11(1):3366. (Clone-specific: Flow cytometry).

Bando JK, Gilfillan S, Di Luccia B, et al. ILC2s are the predominant source of intestinal ILC-derived IL-10. *J Exp Med.* 2020; 217(2):e20191520. (Clone-specific: Flow cytometry).

Patel PS, King RG, Kearney JF. Pulmonary α-1,3-Glucan-Specific IgA-Secreting B Cells Suppress the Development of Cockroach Allergy. *J Immunol.* 2016; 197(8):3175-3187. (Clone-specific: Flow cytometry).

BD Biosciences

bdbiosciences.com

United States 877.232.8995	Canada 888.268.5430	Europe 32.53.720.550	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 0800.771.7157
-------------------------------	------------------------	-------------------------	-----------------------	------------------------------	--

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other

©2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.