

Technical Data Sheet

Mouse IFN- γ ELISPOT Pair

Product Information

Material Number:	551881
Size:	5 plates
Component:	51-2525KZ
Description:	Mouse IFN- γ ELISPOT Capture Antibody
Storage Buffer:	No azide/low endotoxin: Aqueous buffered solution containing no preservative, 0.2 μ m filtered. Endotoxin level is ≤ 0.01 ng/ μ g of protein.
Component:	51-1818KA
Description:	Mouse IFN- γ ELISPOT Detection Antibody
Storage Buffer:	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

Description

The enzyme-linked immunospot (ELISPOT) assay is a powerful tool for detecting and enumerating individual cells that secrete a particular protein in vitro. Based on the sandwich ELISA, the ELISPOT assay derives its specificity and sensitivity by employing high affinity capture and detection antibodies and enzyme-amplification. Although originally developed for analyzing specific antibody-secreting cells, the assay has been adapted for measuring the frequencies of cells that produce and secrete other effector molecules, such as cytokines. The sensitivity of the assay lends itself to measurement of even very low frequencies of cytokine producing cells (e.g., 1/300,000). Unique strengths of the assay include high sensitivity, high throughput, high content analysis, minimal volume of biological material required, applicability to frozen/thawed biological samples, and compatibility with other assays. For example, cells analyzed by ELISPOT can be transferred for cloning, proliferation assays, flow cytometry, or other methods of analysis.

This product contains sufficient reagents for five 96-well plates, including unlabelled anti-cytokine capture antibody (no azide/low endotoxin format), biotinylated anti-cytokine detection antibody and a Certificate of Analysis that provides lot-specific optimal reagent concentrations.

Preparation and Storage

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Recommended Assay Procedure:

ADDITIONAL REAGENTS NEEDED

- BD™ ELISPOT Streptavidin Horseradish Peroxidase (Cat. No. 557630) is recommended.
- Coating Buffer (1X Phosphate Buffered Saline [PBS]): 8 g NaCl; 0.2 g KCl; 1.44 g Na₂HPO₄·7H₂O, 0.24 g KH₂PO₄; dissolved in H₂O to a final volume of 1 liter. Adjust pH to 7.2, sterile-filter (0.2 μ m-sized pore filter) and store at 4°C.
- Blocking Solution: complete tissue culture medium (e.g. RPMI 1640 containing 10% Fetal Bovine Serum [FBS] and 1% Penicillin-Streptomycin-L-Glutamine [Gibco-BRL No. 10378-016]).
- Wash Buffer I: 1X PBS containing 0.05% Tween-20 (0.5 mL Tween-20 per 1 L PBS).
- Wash Buffer II: 1X PBS
- Dilution Buffer: 1X PBS containing 10% FBS.
- BD™ ELISPOT AEC Substrate Set (Cat. No. 551951) is recommended. Alternatively, Prepare as follows:
 - Prepare AEC (3-Amino-9-ethyl-carbazole; Sigma A-5754) stock solution: mix 100 mg AEC in 10 mL DMF (N,N-Dimethylformamide; Sigma D-4551). Caution: dispense DMF in fume hood. Store solution in glassware.
 - Prepare 0.1 M Acetate Solution: add 148 mL of 0.2 M acetic acid/glacial acetic acid to 352 mL of 0.2 M sodium acetate. Adjust volume to 1 L DI water; adjust pH to 5.0.
 - For Final Substrate Solution, add 333.3 μ L of AEC stock solution to 10 mL 0.1 M Acetate Solution. Filter through 0.45 μ m filter. Add 5 μ L of H 202 (30%) and use immediately.

ELISPOT PROTOCOL

NOTE: Use ELISPOT plates and reagents under aseptic conditions (e.g., using a laminar flow hood or biosafety cabinet) for Steps 1-7. Solutions noted with an asterisk (*) are not included and are described in the Additional Reagents Needed section.

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Coating Antibody

1. Dilute capture antibody in Coating Buffer* (see Certificate of Analysis for antibody dilution information). Add 100 µl of diluted antibody solution to each well of an ELISPOT plate; Millipore plate, Cat. No. S2EM004M99 is recommended.
2. Store plates at 4°C overnight.

Blocking

3. Discard Coating Antibody. Wash wells 1X with 200 µL/well Blocking Solution*.
4. Add 200 µL/well Blocking Solution and incubate for 2 hr at room temperature.

Cell Activation

5. Discard Blocking Solution. Prepare mitogen or antigen, diluted in complete tissue culture medium (e.g., RPMI 1640 with FBS, Pen/Strep, and L-glutamine). Add 100 µL/well to ELISPOT plate.
6. Prepare cell suspensions at different densities, (e.g., 1 X 10⁵/mL - 2 X 10⁶ cells/mL) Add 100 µL/well of each cell suspension to ELISPOT plate wells.
7. Replace ELISPOT plate lid. Incubate ELISPOT plate at 37°C, 5% CO₂ and 99% humidity. The duration of the incubation time can be varied (e.g., 2 hr to 24 hr) depending on the nature of the stimulatory cell culture system.

Specific activation conditions and incubation times will vary, depending on cell type, kinetics, and cytokine of interest. Please see Certificate of Analysis provided with each ELISPOT Pair for assay conditions of suggested positive controls.

Note: After step 7, aseptic conditions are no longer needed.

Detection Antibody

8. Aspirate cell suspension. Wash wells 2X with deionized (DI) water. Allow wells to soak for 3-5 min at each wash step.
9. Wash wells 3X with 200 µL/well Wash Buffer I*. Discard Wash Buffer.
10. Dilute Detection Antibody in Dilution Buffer* (see Certificate of Analysis for antibody dilution information). Add 100 µL per well.
11. Replace lid and incubate for 2 hr at room temperature.

Enzyme

12. Discard Detection Antibody solution. Wash wells 3X with 200 µL/well Wash Buffer I*.
13. Dilute Streptavidin-Horseradish Peroxidase (HRP) in Dilution Buffer. (BD™ ELISPOT Streptavidin-HRP, Cat. No. 557630 is recommended). Add 100 µL/well diluted Streptavidin-HRP.
14. Replace lid; incubate for 1 hr at room temperature.

Substrate

15. Discard Streptavidin-HRP solution. Wash wells 4X with 200 µL/well Wash Buffer I*.
16. Wash wells 2X with 200 µL/well Wash Buffer II*.
17. Add 100 µL of Final Substrate Solution (AEC)* to each well. Monitor spot development from 5 - 60 min.
18. Stop substrate reaction by washing wells with DI water.
19. Air-dry plate for 2 hr - overnight at room temperature in the dark, until the plate is completely dry. Store plate in the dark, prior to analysis.
20. Enumerate spots manually using a dissecting microscope or automatically using an ELISPOT Analyzer.

Suggested Companion Products

Catalog Number	Name	Size	Clone
551951	AEC Substrate Set	10 plates	(none)
557630	HRP Streptavidin for ELISPOT	1.0 ml	(none)

Product Notices

1. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Czerkinsky C, Andersson G, Ekre HP, Nilsson LA, Klareskog L, Ouchterlony O. Reverse ELISPOT assay for clonal analysis of cytokine production. I. Enumeration of gamma-interferon-secreting cells. *J Immunol Methods*. 1988; 110(1):29-36.(Methodology)

Fujihashi K, McGhee JR, Beagley KW, et al. Cytokine-specific ELISPOT assay. Single cell analysis of IL-2, IL-4 and IL-6 producing cells. *J Immunol Methods*. 1993; 160(2):181-189.(Methodology)

Helms T, Boehm BO, Asaad RJ, Trezza RP, Lehmann PV, Tary-Lehmann M. Direct visualization of cytokine-producing recall antigen-specific CD4 memory T cells in healthy individuals and HIV patients. *J Immunol*. 2000; 164(7):3723-3732.(Methodology)

Ronnblom L, Cederblad B, Sandberg K, Alm GV. Determination of herpes simplex virus-induced alpha interferon-secreting human blood leucocytes by a filter immuno-plaque assay. *Scand J Immunol*. 1988; 27(2):165-170.(Methodology)

Sedgwick JD, Holt PG. A solid-phase immunoenzymatic technique for the enumeration of specific antibody-secreting cells. *J Immunol Methods*. 1983; 57(1-3):301-309.(Methodology)