## **Technical Data Sheet**

# BUV395 Rat Anti-Mouse Ig, λ1, λ2 & λ3 Light Chain

#### **Product Information**

 Material Number:
 744529

 Size:
 50 μg

 Clone:
 R26-46

Alternative Name: Iql; Iq  $\lambda$ ; Immunoglobulin lambda chain complex; Iq  $\lambda 1/\lambda 2/\lambda 3$ 

Reactivity: Tested in Development:Mouse

 $\begin{tabular}{lll} Isotype: & Rat IgG2a, \kappa \\ Immunogen: & Pooled Mouse Ig \\ \end{tabular}$ 

Application: Flow cytometry(Qualified)

Concentration: 0.2 mg/ml Entrez Gene ID: 111519

Storage Buffer: Aqueous buffered solution containing ≤0.09% sodium azide.

Regulatory Status: RUO

## **Description**

The R26-46 antibody reacts specifically with mouse Igs bearing  $\lambda 1$ ,  $\lambda 2$ , or  $\lambda 3$  light chains. It does not react with  $\kappa$  light chain or heavy chain. Detection of surface immunoglobulin on Ig  $\lambda$  chain-secreting hybridoma cells has been demonstrated with R26-46 mAb.

The antibody was conjugated to BD Horizon™ BUV395 which is part of the BD Horizon Brilliant™ Ultraviolet family of dyes. This dye has been exclusively developed by BD Biosciences to have minimal spillover into other detectors, making it an optimal choice for multicolor flow cytometry. With an Ex Max at 348 nm and an Em Max at 395 nm, BD Horizon BUV395 can be excited with a 355 nm laser and detected with a 379/28 filter.

## **Preparation and Storage**

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with BD Horizon BUV395 under optimal conditions that minimize unconjugated dye and antibody.

# **Recommended Assay Procedure**

For optimal and reproducible results, BD Horizon Brilliant Stain Buffer should be used anytime two or more BD Horizon Brilliant dyes (including BD OptiBuild Brilliant reagents) are used in the same experiment. Fluorescent dye interactions may cause staining artifacts which may affect data interpretation. The BD Horizon Brilliant Stain Buffer was designed to minimize these interactions. More information can be found in the Technical Data Sheet of the BD Horizon Brilliant Stain Buffer (Cat. No. 563794).

## **Suggested Companion Products**

Catalog Number	Name	Size	Clone
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2
554656	Stain Buffer (FBS)	500 mL	
554657	Stain Buffer (BSA)	500 mL	
563794	Brilliant Stain Buffer	100 Tests	
555899	Lysing Buffer	100 mL	
563556	BUV395 Rat IgG2a, κ Isotype Control	50 μg	R35-95

#### **Product Notices**

- 1. This antibody was developed for use in flow cytometry.
- 2. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.

744529 Rev. 1 Page 1 of 2

- 3. Researchers should determine the optimal concentration of this reagent for their individual applications.
- 4. An isotype control should be used at the same concentration as the antibody of interest.
- 5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- 7. Please refer to www.bdbiosciences.com/us/s/resources for technical protocols.
- 8. BD Horizon Brilliant Stain Buffer is covered by one or more of the following US patents: 8,110,673; 8,158,444; 8,575,303; 8,354,239.
- 9. BD Horizon Brilliant Ultraviolet 395 is covered by one or more of the following US patents: 8,158,444; 8,575,303; 8,354,239.

## References

Rolink AG, Winkler T, Melchers F, Andersson J. Precursor B cell receptor-dependent B cell proliferation and differentiation does not require the bone marrow or fetal liver environment.. J Exp Med. 2000; 191(1):23-32. (Clone-specific: Flow cytometry).

Li Y, Li H, Weigert M. Autoreactive B cells in the marginal zone that express dual receptors.. J Exp Med. 2002; 195(2):181-8. (Clone-specific: Flow cytometry).

#### **BD Biosciences**

bdbiosciences.com

 United States
 Canada
 Europe
 Japan
 Asia Pacific
 Latin America/Caribbearn

 877.232.8995
 888.268.5430
 32.53.720.550
 0120.8555.90
 65.6861.0633
 0800.771.7157

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other © 2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.

744529 Rev. 1 Page 2 of 2