

Technical Data Sheet

PE Hamster Anti-Rat CD62L

Product Information

Material Number:	551398
Alternate Name:	Sell; selectin L; L-selectin; LECAM-1; LECAM1; LAM-1; Ly-22
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	HRL1
Immunogen:	Rat LECAM-1 human IgG1 fusion protein
Isotype:	Armenian Hamster IgG2, λ 1
Reactivity:	QC Testing: Rat
Storage Buffer:	Aqueous buffered solution containing \leq 0.09% sodium azide.

Description

The HRL1 monoclonal antibody specifically binds to L-selectin (LECAM-1, CD62L), which is detected on a small percentage of thymocytes and on most neutrophils and peripheral lymphocytes. CD62L is a 62-kDa (on neutrophils) or 65-kDa (on lymphocytes) receptor, with lectin-like and epidermal growth factor-like domains, which binds to sialylated oligosaccharide determinants on high endothelial venules (HEV) in peripheral lymph nodes. This member of the selectin adhesion molecule family appears to be required for lymphocyte homing to peripheral lymph nodes and to contribute to neutrophil emigration at inflammatory sites. L-selectin is rapidly shed from lymphocytes and neutrophils upon cell activation. In the mouse, the level of CD62L expression distinguishes naive CD4+T cells from effector/memory T helper cells. HRL1 antibody inhibits the ligand-binding activity of Lselectin in in vitro assays and slows in vivo leukocyte rolling on microvascular endothelium.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

For detection of L-selectin on peripheral blood leukocytes, use of EDTA as anti-coagulant is recommended; heparin should not be used.

Suggested Companion Products

Catalog Number	Name	Size	Clone
553965	PE Hamster IgG2, λ 1 Isotype Control	0.1 mg	Ha4/8
554656	Stain Buffer (FBS)	500 mL	(none)
554657	Stain Buffer (BSA)	500 mL	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/documents/hamster_chart_11x17.pdf.
5. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
6. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Mulligan MS, Miyasaka M, Tamatani T, Jones ML, Ward PA. Requirements for L-selectin in neutrophil-mediated lung injury in rats. *J Immunol.* 1994; 152(2):832-840. (Biology)

Sprent J, Tough DF. Lymphocyte life-span and memory. *Science.* 1994; 265(5177):1395-1400. (Biology)

Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell.* 1994; 76(2):301-314. (Biology)

Tamatani T, Kitamura F, Kuida K, et al. Characterization of rat LECAM-1 (L-selectin) by the use of monoclonal antibodies and evidence for the presence of soluble LECAM-1 in rat sera. *Eur J Immunol.* 1993; 23(9):2181-2188. (Immunogen: Flow cytometry)

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean
877.232.8995 866.979.9408 32.2.400.98.95 0120.8555.90 65.6861.0633 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

551398 Rev. 13

