

Technical Data Sheet

PE Hamster Anti-Mouse CD3e


Product Information

Material Number:	553240
Alternate Name:	CD3ε chain
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	500A2
Immunogen:	Mouse T-cell receptor
Isotype:	Syrian Hamster IgG2, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

This antibody has been reported to react with the 25-kDa ε chain of the T-cell receptor-associated CD3 complex expressed on thymocytes, mature T lymphocytes, and NK-T cells on many mouse stains. Plate-bound and soluble hamster anti-mouse CD3e (clone 500A2) antibody has been reported to activate unprimed T cells in vitro. Activation of a mouse T-cell clone by soluble hamster anti-mouse CD3e (clone 500A2) can be blocked by Fab fragments of anti-CD4 antibody GK1.5, suggesting that the hamster anti-mouse CD3e (clone 500A2) antibody may bind an epitope of CD3e close to a site at which CD4 associates with the T-cell receptor. This hamster mAb to a mouse leukocyte antigen has been reported not to cross-react with rat leukocytes.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



CD3e expression in spleen and thymus. BALB/c splenocytes were simultaneously stained with FITC-conjugated anti-mouse CD4 mAb RM4-5 (Cat. No. 553046/553047, left panels), FITC-conjugated anti-mouse CD8a mAb 53-6.7 (Cat. No. 553030/553031, left panels), and PE-conjugated mAb 500A2 (bottom left panel). BALB/c thymocytes were stained with PE-conjugated 500A2 (bottom right panel) or unstained (top right panel). Flow cytometry was performed on a BD FACScan™ flow cytometry system.

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed by gel filtration chromatography.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

It has been observed that pre-incubation of thymus cell suspensions at 37°C for 2 to 4 hours prior to staining enhances the ability of anti-CD3e and anti- $\alpha\beta$ TCR mAbs to detect the T-cell receptor on immature thymocytes.

Suggested Companion Products

Catalog Number	Name	Size	Clone
550085	PE Hamster IgG2, κ Isotype Control	0.1 mg	B81-3

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/pharmingen/colors.
4. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

- Allison JP, Havran WL, Poenie M, et al. Expression and function of CD3 on murine thymocytes. In: Kappler J, Davis M, ed. *The T-Cell Receptor, UCLA Symposia, 73rd Edition*. Los Angeles: 1988:33-45.(Immunogen)
- Havran WL, Poenie M, Kimura J, Tsien R, Weiss A, Allison JP. Expression and function of the CD3-antigen receptor on murine CD4+8+ thymocytes. *Nature*. 1987; 330(6144):170-173.(Biology)
- Kubo RT, Born W, Kappler JW, Marrack P, Pigeon M. Characterization of a monoclonal antibody which detects all murine alpha beta T cell receptors. *J Immunol*. 1989; 142(8):2736-2742.(Methodology)
- Ortaldo JR, Winkler-Pickett R, Mason AT, Mason LH. The Ly-49 family: regulation of cytotoxicity and cytokine production in murine CD3+ cells. *J Immunol*. 1998; 160(1):1158-1165.(Biology)
- Portoles P, Rojo J, Golby A, et al. Monoclonal antibodies to murine CD3 epsilon define distinct epitopes, one of which may interact with CD4 during T cell activation. *J Immunol*. 1989; 142(12):4169-4175.(Biology)