

Technical Data Sheet


FITC Rat Anti-Mouse V α 8.3 TCR

Product Information

Material Number:	553376
Size:	0.25 mg
Concentration:	0.5 mg/ml
Clone:	B21.14
Immunogen:	Soluble $\alpha\beta$ TCR from mouse cytotoxic T-cell clone CW3/1.1
Isotype:	Rat (LOU) IgG1, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The B21.14 antibody reacts with some members of the V α 8 T-cell Receptor (TCR) subfamily of mice having the *a*, *b*, *c*, and *d* haplotypes of the *Tcr* gene complex (e.g., all strains tested). It recognizes an epitope in the CDR1 of V α 8.3, but not V α 8.2, TCR subfamily member, as does the KT50 mAb (Cat. No. 557096, for the biotinylated antibody). Site-directed mutagenesis has identified three amino acids which are necessary for antibody reactivity and which are unique to V α 8.3 among the five functional V α 8 TCR subfamily members. On a common H-2K[k] background and using the KT50 mAb, the frequency of V α 8.3 TCR-bearing T lymphocytes has been demonstrated to be higher in *Tcr*[*a/a*] mice than in *Tcr*[*a/b*] mice. Furthermore, studies of congenic strains suggest that CD8⁺ V α 8.3 TCR-bearing T lymphocytes undergo negative selection in mice expressing MHC class I antigens of the *H-2[d]* haplotype.



Two-color analysis of the expression of Va 8.3 TCR on peripheral lymphocytes. C57BL/6 lymph node cells were incubated simultaneously with FITC-conjugated B21.14, PE-conjugated anti-mouse CD4 RM4-5 (Cat. No. 553048/553049), and PE-conjugated anti-mouse CD8a 53-6.7 (Cat. No. 553032/553033) monoclonal antibodies. Flow cytometry was performed on a BD FACScan™ flow cytometry system.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry

Routinely Tested

Recommended Assay Procedure:

For flow cytometry of cell suspensions from peripheral lymphoid tissues, it is recommended that multicolor staining be performed to distinguish T lymphocytes from non-T cells.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
553048	PE Rat Anti-Mouse CD4	0.1 mg	RM4-5
553032	PE Rat Anti-Mouse CD8a	0.1 mg	53-6.7
553924	FITC Rat IgG1, κ Isotype Control	0.25 mg	R3-34

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

- Brodnicki TC, Holman PO, Kranz DM. Reactivity and epitope mapping of single-chain T cell receptors with monoclonal antibodies. *Mol Immunol.* 1996; 33(3):253-263.(Biology)
- Necker A, Rebai N, Matthes M, et al. Monoclonal antibodies raised against engineered soluble mouse T cell receptors and specific for V alpha 8-, V beta 2- or V beta 10-bearing T cells. *Eur J Immunol.* 1991; 21(12):3035-3040.(Immunogen)
- Tomonari K. Negative selection of Tcr α -V8+CD8+ T cells by MHC class I molecules. *Immunogenetics.* 1992; 35(5):291-295.(Biology)
- Tomonari K, Fairchild S, Rosenwasser OA. Influence of viral superantigens on V beta- and V alpha-specific positive and negative selection. *Immunol Rev.* 1993; 131:131-168.(Biology)