

Technical Data Sheet

Perm/Wash Buffer I

Product Information

Material Number:	557885
Size:	125 mL
Storage Buffer:	Aqueous buffered solution containing saponin, fetal bovine serum and $\leq 0.09\%$ sodium azide.

Description

BD Phosflow™ Perm/Wash Buffer I is intended to be used for the intracellular staining of post-translationally modified signaling proteins. BD Phosflow™ Perm/Wash Buffer I is used to permeabilize cells and to serve as an antibody diluent and cell wash buffer. It is optimized for use with the BD Phosflow™ brand of intracellular phosphorylated signaling protein-specific antibodies. BD Phosflow™ Perm/Wash Buffer I is provided as a 10X concentrated solution containing FBS and saponin. The presence of a small amount of precipitate may be observable and will not affect the performance of the buffer. Because saponin-mediated cell permeabilization is a reversible process, it is important to keep the cells in the presence of saponin during intracellular staining.

Preparation and Storage

Store undiluted at 4°C.

Do not freeze.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

Flow cytometry: Dilute BD Phosflow™ Perm/Wash Buffer I 1:10 in distilled water prior to use. If desired, the diluted 1X BD™ Phosflow Perm/Wash Buffer I can be passed through a 0.45 µm filter to remove any residual precipitate. The BD Phosflow™ Perm/Wash Buffer I can be used to permeabilize or wash cells and to dilute antibodies for immunofluorescent staining of intracellular proteins.

Suggested Companion Products

Catalog Number	Name	Size	Clone
557870	Fix Buffer I	250 mL	(none)
558049	Lyse/Fix Buffer 5X	250 mL	(none)
554655	Fixation Buffer	100 mL	(none)

Product Notices

- Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Assenmacher M, Schmitz J, Radbruch A. Flow cytometric determination of cytokines in activated murine T helper lymphocytes: expression of interleukin-10 in interferon-gamma and in interleukin-4-expressing cells. *Eur J Immunol.* 1994; 24(5):1097-1101. (Biology)

Elson LH, Nutman TB, Metcalfe DD, Prussin C. Flow cytometric analysis for cytokine production identifies T helper 1, T helper 2, and T helper 0 cells within the human CD4+CD27- lymphocyte subpopulation. *J Immunol.* 1995; 154(9):4294-4301. (Biology)

Jung T, Schauer U, Heusser C, Neumann C, Rieger C. Detection of intracellular cytokines by flow cytometry. *J Immunol Methods.* 1993; 159(1-2):197-207. (Biology)

Krutzik PO, Nolan GP. Intracellular phospho-protein staining techniques for flow cytometry: monitoring single cell signaling events. *Cytometry A.* 2003; 55(2):61-70. (Biology)

Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods.* 1995; 188(1):117-128. (Biology)

Sander B, Andersson J, Andersson U. Assessment of cytokines by immunofluorescence and the paraformaldehyde-saponin procedure. *Immunol Rev.* 1991; 119:65-93. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

