

Technical Data Sheet


Perm Buffer II

Product Information

Material Number:	558052
Size:	125 mL
Storage Buffer:	Aqueous buffered solution containing proprietary ingredients.

Description

BD Phosflow™ Perm Buffer II is intended to be used for the permeabilization of fixed PBMC's or lysed whole blood. PBMC's, for example, can be stimulated with kinase activators, such as phorbol esters, and can be fixed with BD Phosflow™ Fix Buffer I (Cat. No. 557870) followed by permeabilization of the cells with BD Phosflow™ Perm Buffer II to allow for an effective intracellular staining of kinases. Designed for optimal fluorescent staining in flow cytometry, BD Phosflow™ Perm Buffer II can be used with antibodies for phospho-protein analysis (i.e the staining of intracellular proteins involved in cell signalling pathways, such as phosphorylated proteins and assorted kinases). In many cases, these phospho-specific antibodies may be used simultaneously with the staining of cell surface markers (e.g CD markers). Not all cell surface markers, however, are compatible with BD Phosflow™ Perm Buffer II. For a current listing of compatible cell surface markers for use with BD Phosflow™ Perm Buffer II, investigators are encouraged to visit the following website: http://wwwbdbiosciences.com/documents/antibodies_human_cellsurface_marker.pdf.



Flow cytometric analysis of ERK1/2 (pT202/pY204) in human PBMC. Freshly isolated human PBMC were either left untreated (unshaded histogram) or treated (shaded histogram) with 40 nM PMA at 37°C for 10 minutes. The PBMC were then fixed using BD Phosflow™ Fix Buffer I (Cat.No. 557870) and permeabilized with BD Phosflow™ Perm Buffer II (Cat. No. 558052) for 30 minutes on ice. After washing the cells twice with BD Pharmingen™ Stain Buffer (Cat. No. 554656/554657), the PBMC were stained with Alexa Fluor® 488 Rat Anti-ERK1/2 (pT202/pY204) (Cat. No. 512592).

Preparation and Storage

Store undiluted at room temperature.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

Flow cytometry:

1. Warm BD Phosflow™ Fix Buffer I (Cat. No. 557870) to 37 °C prior to use.
2. Chill BD Phosflow™ Perm Buffer II to -20 °C prior to use.
3. Collect PBMCs after Ficoll-Paque separation and treat with any appropriate cell activators as applicable.
4. At the end of the treatment, immediately mix one volume of the warmed BD Phosflow™ Fix Buffer I with one volume of the PBMC suspension. Mix well and incubate the tubes in a 37 °C water bath for 10 min. Spin down the cells at 250 x g for 10 min in a table-top centrifuge and then aspirate the supernatant.
5. Wash the cells once with BD Pharmingen™ Stain Buffer (FBS) (Cat. No. 554656). Alternatively, 1X Washing/Staining Solution (1 x PBS, 1% FCS, and 0.09% sodium azide) may be used. Centrifuge to pellet the cells and remove the supernatant.
6. Vortex to loosen the cells. Permeabilize the cells by slowly adding cold BD Phosflow™ Perm Buffer II while vortexing. Incubate on ice for 30 min and then spin down the cells.
7. Wash the cells twice with 5-10 mL of BD Pharmingen™ Stain Buffer (FBS). Centrifuge at 250 x g for 10 minutes and remove the supernatant by aspiration.
8. Resuspend cells in BD Pharmingen™ Stain Buffer (FBS) at 1x 10⁷/ml and aliquot 100 µl to each flow tube to continue with antibody staining and flow cytometric analysis.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 866.979.9408 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.



Danger: BD Phosflow™ Perm Buffer II contains 64.86% methanol (w/w).

Hazard statements

Highly flammable liquid and vapor.

Toxic if swallowed, in contact with skin or if inhaled.

Causes damage to organs.

Precautionary statements

Keep away from heat/sparks/open flames/hot surfaces/ignition sources. - No smoking.

Keep container tightly closed. Ground and bond container and receiving equipment. Use explosion-proof [electrical/ventilating/lighting/] equipment.

Use non-sparking tools. Take action to prevent static discharges.

Wear protective clothing/eye protection/gloves.

Do not breathe mist/vapours/spray.

Use only outdoors or in a well-ventilated area. Wash thoroughly after handling. Do not eat, drink or smoke when using this product.

IF ON SKIN (or hair): Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Take off immediately all contaminated clothing. Wash contaminated clothing before reuse.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

IF SWALLOWED: Immediately call a POISON CENTER/doctor. Rinse mouth. Call a POISON CENTER/doctor.

If exposed: Call a poison center/doctor

Store in well-ventilated place. Keep cool. Store locked up. Keep container tightly closed.

Static accumulating flammable liquid can become electrostatically charged even in bonded and grounded equipment.

Sparks may ignite liquid and vapor. May cause flash fire or explosion.

Use suitable extinguishing media to extinguish

Suggested Companion Products

Catalog Number	Name	Size	Clone
557870	Fix Buffer I	250 mL	(none)
554656	Stain Buffer (FBS)	500 mL	(none)
612592	Alexa Fluor® 488 Mouse Anti-ERK1/2 (pT202/pY204)	50 Tests	20A
554655	Fixation Buffer	100 mL	(none)
558049	Lyse/Fix Buffer 5X	250 mL	(none)
557885	Perm/Wash Buffer I	125 mL	(none)
558050	Perm Buffer III	125 mL	(none)
554657	Stain Buffer (BSA)	500 mL	(none)

Product Notices

1. Ficoll-Paque is a trademark of Amersham Biosciences Limited.
2. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
4. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
5. Please refer to www.bdbiosciences.com/us/s/resources for technical protocols.