

Technical Data Sheet

Purified Mouse Anti-Rat CD26

Product Information

Material Number:	559639
Alternate Name:	Dpp4; dipeptidyl peptidase IV; DPPIV; Cd26
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	OX-61
Immunogen:	Rat dendritic cells enriched from thoracic duct lymph
Isotype:	Mouse (BALB/c) IgG2a, κ
Reactivity:	QC Testing: Rat
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The OX-61 monoclonal antibody specifically recognizes a type II transmembrane glycoprotein, CD26, which is a serine exoprotease identified as dipeptidyl/peptidase IV. Among various biological activities, rat CD26 binds fibronectin and collagen. Rat CD26 is involved in the costimulation of thymocyte proliferation in vitro, particularly the CD4-/CD8- subset, and is developmentally regulated on hematopoietic cells. Although mouse and human CD26 anchor ADA (adenosine deaminase) to cell membranes, rat CD26 does not function as an ADA-binding protein. Rat CD26 is expressed in lung endothelial cells, as well as in various epithelial cells. T cells express lower levels of CD26 than CD4-/CD8- thymocytes. The distribution of CD26 antigen in rat bone marrow cells is similar to that of human CD26. OX-61 monoclonal antibody stains CD4+, CD8+, and Ig+ lymphocytes, and the staining increases upon activation.

Preparation and Storage

Store undiluted at 4°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Reported
Western blot	Reported
Immunoprecipitation	Reported

Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
553454	Purified Mouse IgG2a κ Isotype Control	0.5 mg	G155-178
554656	Stain Buffer (FBS)	500 mL	(none)
554657	Stain Buffer (BSA)	500 mL	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE (No Azide/Low Endotoxin) antibody format, if available, for in vitro and in vivo use.
5. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Bristol LA, Sakaguchi K, Appella E, Doyle D, Takacs L. Thymocyte costimulating antigen is CD26 (dipeptidyl-peptidase IV). Costimulation of granulocyte, macrophage, and T lineage cell proliferation via CD26. *J Immunol.* 1992; 149(2):367-372. (Biology)

Gorrell MD, Wickson J, McCaughan GW. Expression of the rat CD26 antigen (dipeptidyl peptidase IV) on subpopulations of rat lymphocytes. *Cell Immunol.* 1991; 134(1):205-215. (Clone-specific: Immunohistochemistry, Immunoprecipitation)

Hanski C, Huhle T, Gossrau R, Reutter W. Direct evidence for the binding of rat liver DPP IV to collagen in vitro. *Exp Cell Res.* 1988; 178(1):64-72. (Biology)

Iwaki-Egawa S, Watanabe Y, Fujimoto Y. CD26/dipeptidyl peptidase IV does not work as an adenosine deaminase-binding protein in rat cells. *Cell Immunol.* 1997; 178(2):180-186. (Biology)

Johnson RC, Zhu D, Augustin-Voss HG, Pauli BU. Lung endothelial dipeptidyl peptidase IV is an adhesion molecule for lung-metastatic rat breast and prostate carcinoma cells. *J Cell Biol.* 1993; 121(6):1423-1432. (Biology)

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean
877.232.8995 866.979.9408 32.2.400.98.95 0120.8555.90 65.6861.0633 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

559639 Rev. 5



Kameoka J, Tanaka T, Nojima Y, Schlossman SF, Morimoto C. Direct association of adenosine deaminase with a T cell activation antigen, CD26. *Science*. 1993; 261(5120):466-469. (Biology)

McCaughan GW, Wickson JE, Creswick PF, Gorrell MD. Identification of the bile canalicular cell surface molecule GP110 as the ectopeptidase dipeptidyl peptidase IV: an analysis by tissue distribution, purification and N-terminal amino acid sequence. *Hepatology*. 1990; 11(4):534-544. (Immunogen: Immunohistochemistry, Immunoprecipitation, Western blot)

Piazza GA, Callanan HM, Mowery J, Hixson DC. Evidence for a role of dipeptidyl peptidase IV in fibronectin-mediated interactions of hepatocytes with extracellular matrix. *Biochem J*. 1989; 262(1):327-334. (Biology)