

Technical Data Sheet


Alexa Fluor® 647 Mouse anti-PKA RIIα (pS99)**Product Information**

Material Number:	560164
Size:	50 Tests
Vol. per Test:	20 µl
Clone:	I65-856.286
Immunogen:	Phosphorylated Human PKA RIIα Peptide
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

cAMP-dependent Protein Kinase (PKA) is composed of two distinct subunits: catalytic (C) and regulatory (R). Four regulatory subunits have been identified: RIα, RIβ, RIIα, and RIIβ. These subunits define type I and II PKAs. Following binding of cAMP, the regulatory subunits dissociate from the catalytic subunits, rendering the enzyme active. Type I and II holoenzymes have three potential C subunits (Cα, Cβ, or Cγ). Type II PKA can be distinguished by autophosphorylation of the R subunits, while type I PKA binds Mg/ATP with high affinity. The cAMP-dependent autophosphorylation of the human RIIα subunit occurs at Serine 99 (S99) [Entrez Protein Accession #CAA33094]. Most cells express both type I and type II PKAs. Although the Rα isoforms are ubiquitously expressed, the Rβ isoforms are predominantly found in nervous and adipose tissues. In addition to their enzyme regulatory activity, the RIIα and RIIβ subunits determine the subcellular location of the holoenzymes via their interactions with specific intracellular anchoring proteins.

The I65-856.286 antibody recognizes human PKA RIIα phosphorylated at S99.



Analysis of PKA RIIα (pS99) in human peripheral blood lymphocytes. Human peripheral blood mononuclear cells (PBMC) were either treated with 1 µM Staurosporine (EMD Biosciences, Cat. No. 569397) for 2 hours at 37°C (shaded histogram) or untreated (open histogram). The PBMC were fixed (BD Cytotfix™ buffer, Cat. No. 554655) for 10 minutes at 37°C, permeabilized with BD Phosflow™ Perm Buffer III (Cat. No. 558050) on ice for 30 minutes, and then stained with Alexa Fluor® 647 Mouse anti-PKA RIIα (pS99, Cat. No. 560164). For data analysis, lymphocytes were selected by their scatter profile. The data demonstrates that the level of phosphorylation of PKA RIIα decreases when protein kinase activity is inhibited by the treatment. Flow cytometry was performed on a BD FACSCalibur™ flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Application Notes**Application**

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

Either BD Cytotfix™ fixation buffer or BD Phosflow™ Fix Buffer I may be used for cell fixation. Any of the three BD Phosflow™ permeabilization buffers may be used.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 866.979.9408 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2016 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

560164 Rev. 2



The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	Daudi	Staurosporine	Cytofix	Perm III	Unsatisfactory
	Human	EA-hy 926	Staurosporine	Cytofix	Perm III	Down-regulation
	Human	HeLa S3	Calyculin A + Okadaic Acid	Cytofix	Perm III	Unsatisfactory
	Human	Jurkat	Staurosporine	Cytofix	Perm III	Down-regulation
	Human	PBMC	Staurosporine	Cytofix	Perm I, II, or III	Down-regulation
WB	Human	Daudi	Staurosporine			no band
	Human	EA-hy 926	Staurosporine			51-kDa band down-regulated
	Human	EA-hy 926	Calyculin A + Okadaic Acid			no induction
	Human	HeLa S3	Calyculin A + Okadaic Acid			51-kDa band induced
	Human	Jurkat	Staurosporine			51-kDa band down-regulated
	Human	PBMC	Staurosporine			51-kDa band down-regulated

Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 mL	(none)
557870	Fix Buffer I	250 mL	(none)
558050	Perm Buffer III	125 mL	(none)
558052	Perm Buffer II	125 mL	(none)
557885	Perm/Wash Buffer I	125 mL	(none)
554656	Stain Buffer (FBS)	500 mL	(none)
554657	Stain Buffer (BSA)	500 mL	(none)

Product Notices

- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
- Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
- Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Francis SH, Corbin JD. Structure and function of cyclic nucleotide-dependent protein kinases. *Annu Rev Physiol.* 1994; 56:237-272. (Biology)
 Scott JD, Stofko RE, McDonald JR, Comer JD, Vitalis EA, Magili JA. Type II regulatory subunit dimerization determines the subcellular localization of the cAMP-dependent protein kinase. *J Biol Chem.* 1990; 35:21561-21566. (Biology)

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 866.979.9408 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2016 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

560164 Rev. 2

