

## Technical Data Sheet


**Pacific Blue™ Mouse anti-Stat1 (pY701)****Product Information**

<b>Material Number:</b>	<b>560310</b>
<b>Size:</b>	50 Tests
<b>Vol. per Test:</b>	20 µl
<b>Clone:</b>	14/P-STAT1
<b>Immunogen:</b>	Phosphorylated Human Stat1 (pY701) Peptide
<b>Isotype:</b>	Mouse IgG1, κ
<b>Reactivity:</b>	QC Testing: Human Tested in Development: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09% sodium azide.

**Description**

Stat (Signal transducer and activators of transcription) proteins are critical mediators of the biologic activity of cytokines, including interleukins, interferons, erythropoietin, and growth factors. Ligand-receptor interaction leads to activation of constitutively associated JAK family kinases and subsequent recruitment/activation of Stat proteins by tyrosine phosphorylation. Active Stat proteins then move to the nucleus to promote transcription of cytokine-inducible genes. Seven Stat proteins have been cloned, each of which is differentially expressed and/or activated in a cytokine-specific and cell type-specific manner. Stat1 and Stat2 are components of the ISGF3 (Interferon-Stimulated Gene Factor 3) complex, which is the primary transcription activator induced by the binding of the interferon to a specific cell-surface receptor. Stat1 has two alternatively spliced isoforms, 91-kDa Stat1α and 84-kDa Stat1β; Stat1α has 38 additional C-terminal amino acids. In response to the binding of IFNα, IFNγ, EGF, PDGF, or CSF-1 to their respective receptors, the Stat1 subunits become tyrosine-phosphorylated at Y701, and the complex is translocated to the nucleus. This results in the formation of an active complex that includes the DNA-binding p48 subunit. This complex is responsible for modulating the transcription of the interferon-stimulated genes (ISGs).

The 14/P-STAT1 monoclonal antibody recognizes the phosphorylated Y701 in Stat1α and Stat1β.



**Analysis of Stat1 (pY701) in human peripheral blood monocytes.** Peripheral blood mononuclear cells (PBMC) were either left unstimulated (unshaded) or stimulated (shaded) with 100 ng/ml (final concentration) BD Pharming™ Recombinant Human IFNγ (Cat. No. 554617) for 15 minutes at 37°C. The PBMC were fixed (BD Cytotfix™ Fixation buffer, Cat. No. 554655) for 10 minutes at 37°C, then permeabilized (BD Phosflow™ Perm Buffer III, Cat. No. 558050) on ice for 30 minutes and then stained with Pacific Blue™ Mouse anti-Stat1(pY701). For data analysis, monocytes were selected by scatter profile. Flow cytometry was performed on a BD FACSCanto™ II flow cytometer.

**Preparation and Storage**

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody is conjugated to Pacific Blue™ under optimum conditions, and unreacted Pacific Blue™ was removed.

**BD Biosciences**

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD



The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	PBMC	human IFN $\gamma$	Cytofix	Perm III	Upregulated expression
		PBMC	human IFN $\gamma$	Cytofix	Perm I or II	Unsatisfactory
		U937	human IFN $\gamma$	Cytofix	Perm III	Upregulated expression
		U937	human IFN $\gamma$	Cytofix	Perm I or II	Unsatisfactory
WB	Human	A-431	EGF			84- & 91-kDa bands induced

## Application Notes

### Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554617	Recombinant Human IFN- $\gamma$	50 $\mu$ g	(none)
554655	Fixation Buffer	100 mL	(none)
558050	Perm Buffer III	125 mL	(none)

### Product Notices

- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100- $\mu$ l experimental sample (a test).
- Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
- The Alexa Fluor<sup>®</sup>, Pacific Blue<sup>™</sup>, and Cascade Blue<sup>®</sup> dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor<sup>®</sup> dyes (except for Alexa Fluor<sup>®</sup> 430), Pacific Blue<sup>™</sup> dye, and Cascade Blue<sup>®</sup> dye are covered by pending and issued patents.
- Pacific Blue<sup>™</sup> has a maximum absorption of 416 nm and maximum emission of 451 nm. Before staining with this reagent, please confirm that your flow cytometer is capable of exciting the fluorochrome and discriminating the resulting fluorescence.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
- Species testing during development may have been performed with a different format of the same clone. Selected applications have been tested for cross-reactivity.

### References

Bromberg J, Darnell JE. The role of STATs in transcriptional control and their impact on cellular function. *Oncogene*. 2000; 19(21):2468-2473. (Biology)  
 Darnell JE Jr. STATs and gene regulation. *Science*. 1997; 277(5332):1630-1635. (Biology)  
 Fu XY, Zhang JJ. Transcription factor p91 interacts with the epidermal growth factor receptor and mediates activation of the c-fos gene promoter. *Cell*. 1993; 74(6):1135-1145. (Biology)

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States 877.232.8995    Canada 800.268.5430    Europe 32.2.400.98.95    Japan 0120.8555.90    Asia Pacific 65.6861.0633    Latin America/Caribbean 55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

