

Technical Data Sheet


Alexa Fluor® 488 Mouse anti-Oct3/4 (Human Isoform A)

Product Information


Material Number:	561628
Alternate Name:	Oct3/4A, Oct-3A, OTF-3, NF-A3, OTF4, POU5F1, MGC22487
Entrez Gene ID:	5460, 18999
Size:	50 tests
Vol. per Test:	5 µl
Clone:	O50-808
Immunogen:	Human Oct3/4 Isoform A Recombinant Protein
Isotype:	Mouse (BALB/c) IgG1, κ
Reactivity:	QC Tested: Human
Storage Buffer:	Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09% sodium azide.

Description

Development of a multicellular organism from a single fertilized egg is regulated by the coordinated activity of DNA transcription factors. Oct3/4, a member of the POU family of transcription factors, functions in pluripotent cells of early embryonic stem (ES) cell lines and embryonal carcinomas (EC). The human POU5F1 gene can encode various splice variants, two of which are Oct3/4A and Oct3/4B. Both isoforms share identical POU DNA-binding and C-terminal domains but differ in their N-terminal domain. The N-terminal domain of Oct3/4B is inhibitory to the DNA binding domain and therefore cannot stimulate transcription of Oct3/4-dependent genes. Oct3/4B can be detected in both pluripotent and some differentiated cell types in both the nucleus and cytoplasm, but its function is unclear. There is not an equivalent to Oct3/4B in mouse. Oct3/4A is expressed in the nucleus and has been demonstrated to orchestrate the transcription of Oct3/4-dependent genes. It has been demonstrated that the expression of Oct3/4 isoforms can vary greatly in different cell types, and discrimination of these is crucial for assessing Oct3/4 expression and function. The O50-808 monoclonal antibody recognizes human Oct3/4 Isoform A and mouse Oct3/4.



Analysis of Oct3/4 Isoform A on human embryonic stem (ES) cells. H9 human ES cells (WiCell, Madison, WI) were harvested, fixed in BD Cytotfix™ fixation buffer (Cat. No. 554655), permeabilized with BD Phosflow™ Perm Buffer III (Cat. No. 558050) and stained with matching concentrations of either Alexa Fluor® 488 Mouse IgG1, κ isotype control (Dashed line, Cat. No. 557782) or Alexa Fluor® 488 Mouse anti-Oct3/4 (Human Isoform A) monoclonal antibody (solid line). The histograms were derived from gated events based on light scattering characteristics for the H9 cell line. Flow cytometry was performed on a BD LSR™ II flow cytometry system.



Immunofluorescent staining of Oct3/4 Isoform A in human embryonic stem (ES) cells. H9 human ES cells (WiCell, Madison, WI) passage 33 grown on irradiated mouse fibroblasts were fixed with BD Cytotfix™ fixation buffer (Cat. No. 554655), permeabilized, and stained with Alexa Fluor® 488 Mouse anti-Oct3/4 (Human Isoform A) monoclonal antibody (pseudo-colored green) at 2.5 µg/mL. Cell nuclei were counter stained with Hoechst 33342 (pseudo-colored blue). The images were captured on a BD Pathway™ 435 Cell Analyzer and merged using BD Attovision™ Software. The cells were permeabilized with BD™ Phosflow Perm Buffer III (Cat. No. 558050); 1x BD Perm/Wash™ Buffer (Cat No. 554723) is also suitable for permeabilization.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 488 under optimum conditions, and unreacted Alexa Fluor® 488 was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
Bioimaging	Tested During Development
Immunofluorescence	Tested During Development

Suggested Companion Products

Catalog Number	Name	Size	Clone
557782	Alexa Fluor® 488 Mouse IgG1 κ Isotype Control	50 tests	MOPC-21
554655	Fixation Buffer	100 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)
554723	Perm/Wash Buffer	100 ml	(none)
353219	BD Falcon™ 96-well Imaging Plate	NA	(none)
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100-µl experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. An isotype control should be used at the same concentration as the antibody of interest.
4. Alexa Fluor® 488 fluorochrome emission is collected at the same instrument settings as for fluorescein isothiocyanate (FITC).
5. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
6. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
7. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
8. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
9. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.

References

Nishimoto M, Fukushima A, Okuda A, Muramatsu M. The gene for the embryonic stem cell coactivator UTF1 carries a regulatory element which selectively interacts with a complex composed of Oct-3/4 and Sox-2. *Mol Cell Biol.* 1999; 19(8):5453-5465. (Biology)

Okamoto K, Okazawa H, Okuda A, Sakai M, Muramatsu M, Hamada H. A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. *Cell.* 1990; 60(3):461-472. (Biology)

Pan G, Thomson JA. Nanog and transcriptional networks in embryonic stem cell pluripotency. *Cell Res.* 2007; 17:42-49. (Biology)

Rosfjord E, Scholtz B, Lewis R, Rizzino A. Phosphorylation and DNA binding of the octamer binding transcription factor Oct-3. *Biochem Biophys Res Commun.* 1995; 212(3):847-853. (Biology)

Vigano MA, Staudt LM. Transcriptional activation by Oct-3: evidence for a specific role of the POU-specific domain in mediating functional interaction with Oct-1. *Nucleic Acids Res.* 1996; 24(11):2112-2118. (Biology)

Yuan H, Corbi N, Basilico C, Dailey L. Developmental-specific activity of the FGF-4 enhancer requires the synergistic action of Sox2 and Oct-3. *Genes Dev.* 1995; 9(21):2635-2645. (Biology)