

Technical Data Sheet


APC Rat Anti-Mouse IL-4

Product Information

Material Number:	562045
Size:	25 µg
Concentration:	0.2 mg/ml
Clone:	11B11
Immunogen:	Partially Purified Mouse IL-4
Isotype:	Rat IgG1
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The 11B11 antibody reacts with mouse interleukin-4 (IL-4). The immunogen used to generate the 11B11 hybridoma was partially purified mouse IL-4 from PMA-stimulated EL-4 supernatant. The purified or unconjugated form of this antibody has been reported to be neutralizing.



Expression of IL-4 by stimulated CD4+ and CD4-Balb/c spleen cells. BALB/c spleen cells were cultured for 72 hours in medium containing *Staphylococcus aureus* enterotoxin B (2 µg/ml final concentration; Sigma, St. Louis, MO), recombinant mouse IL-2 (10 U/ml final concentration; Cat. No. 550069) and recombinant mouse IL-4 (2 ng/ml final concentration; Cat. No. 550067). The cells were harvested and restimulated for 5 hours with anti-CD3 (2 µg/ml final concentration; 145-2C11, Cat. No. 553057) and anti-CD28 (2 µg/ml final concentration; clone 37.51, Cat. No. 553294) antibodies in the presence of GolgiStop™ (3 µM final concentration; Cat. No. 554724). The splenocytes were then stained with 0.25 µg of FITC-conjugated rat anti-mouse CD4 (FITC-RM4-5, Cat. No. 553047) and 0.25 µg of APC-conjugated rat anti-mouse IL-4 antibody (APC-11B11, Cat. No. 554436) by using Pharmingen's staining protocol (left panel). To demonstrate staining specificity, the binding of APC-11B11 was blocked by the preincubation of the conjugated antibody with excess recombinant mouse IL-4 (0.25 µg; Cat. No. 550037) (middle panel) or by pre-blocking fixed/permeabilized cells with excess purified "cold" 11B11 mouse antibody (5.0 µg; Cat. No. 554433) (right panel) prior to staining. The quadrant markers for the bivariate dot plots were set based on the autofluorescence control, and verified using the cytokine-blocking and "cold" mouse antibody blocking controls. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe or red diode laser. These include the dual laser FACStarPLUS™, FACS Vantage™ or FACSCalibur™.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Application Notes

Application

Intracellular staining (flow cytometry)

Routinely Tested

Recommended Assay Procedure:

Immunofluorescent Staining and Flow Cytometric Analysis: The APC-conjugated 11B11 antibody is useful for immunofluorescent staining and flow cytometric analysis to identify and enumerate IL-4 producing cells within mixed cell populations. The use of an isotype control, such as rat IgG1 isotype control, APC-R3-34 (Cat. No. 554686) is recommended. For specific methodology, please visit our web site, www.bdbiosciences.com, and go to the Technical Resources/protocols section or the chapter on intracellular staining in the Immune Function Handbook.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
550067	Recombinant Mouse IL-4	10 µg	(none)
554686	APC Rat IgG1, κ Isotype Control	0.1 mg	R3-34
550069	Recombinant Mouse IL-2	20 µg	(none)
553057	Purified NA/LE Hamster Anti-Mouse CD3e	0.5 mg	145-2C11
553294	Purified NA/LE Hamster Anti-Mouse CD28	0.5 mg	37.51
554724	Protein Transport Inhibitor (Containing Monensin)	0.7 ml	(none)
554653	MiCK-2 Mouse Cytokine Positive Control Cells	1.0 ml	(none)
554715	BD Cytotfix/Cytoperm Plus Kit (with BD GolgiStop)	250 tests	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to wwwbdbiosciences.com/pharming/en/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at wwwbdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.
6. An isotype control should be used at the same concentration as the antibody of interest.

References

Abrams J. Immunoenzymetric assay of mouse and human cytokines using NIP-labeled anti-cytokine antibodies. In: Coligan J, Kruisbeek A, Margulies D, Shevach E, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley and Sons; 1995:6.20-6.21. (Clone-specific: ELISA)

Assenmacher M, Schmitz J, Radbruch A. Flow cytometric determination of cytokines in activated murine T helper lymphocytes: expression of interleukin-10 in interferon-gamma and in interleukin-4-expressing cells. *Eur J Immunol*. 1994; 24(5):1097-1101. (Clone-specific: Flow cytometry)

Haak-Frendscho M, Brown JF, Iizawa Y, Wagner RD, Czuprynski CJ. Administration of anti-IL-4 monoclonal antibody 11B11 increases the resistance of mice to *Listeria monocytogenes* infection. *J Immunol*. 1992; 148(12):3978-3985. (Clone-specific: Neutralization)

Lindqvist C, Lundstrom H, Oker-Blom C, Akerman KE. Enhanced IL-4-mediated D10.G4.1 proliferation with suboptimal concentrations of anti-IL-4 receptor monoclonal antibodies. *J Immunol*. 1993; 150(2):394-398. (Clone-specific: Neutralization)

Ohara J, Paul WE. Production of a monoclonal antibody to and molecular characterization of B-cell stimulatory factor-1. *Nature*. 1985; 315(6017):333-336. (Immunogen)

Openshaw P, Murphy EE, Hosken NA, et al. Heterogeneity of intracellular cytokine synthesis at the single-cell level in polarized T helper 1 and T helper 2 populations. *J Exp Med*. 1995; 182(5):1357-1367. (Clone-specific: Flow cytometry)

Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods*. 1995; 188(1):117-128. (Methodology)

Sadick MD, Heinzel FP, Holaday BJ, Pu RT, Dawkins RS, Locksley RM. Cure of murine leishmaniasis with anti-interleukin 4 monoclonal antibody. Evidence for a T cell-dependent, interferon gamma-independent mechanism. *J Exp Med*. 1990; 171(1):115-127. (Clone-specific: Neutralization)

Sander B, Hoiden I, Andersson U, Moller E, Abrams JS. Similar frequencies and kinetics of cytokine producing cells in murine peripheral blood and spleen. Cytokine detection by immunoassay and intracellular immunostaining. *J Immunol Methods*. 1993; 166(2):201-214. (Clone-specific: ELISA, Flow cytometry)

Swain SL, Weinberg AD, English M, Huston G. IL-4 directs the development of Th2-like helper effectors. *J Immunol*. 1990; 145(11):3796-3806. (Clone-specific: Neutralization)