


## Technical Data Sheet

**PE Rat anti-Mouse CD370 (Clec9A)****Product Information**

<b>Material Number:</b>	<b>562734</b>
<b>Alternate Name:</b>	CD370; CLC9A; Clec9a; DNGR-1; C-type lectin domain family 9 member A
<b>Size:</b>	50 µg
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	10B4 (also known as 24/04-10B4)
<b>Immunogen:</b>	Mouse Clec9A Peptide
<b>Isotype:</b>	Rat (WI) IgG2a, κ
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing ≤0.09% sodium azide.

**Description**

The 10B4 monoclonal antibody specifically binds to mouse Clec9A. Mouse Clec9A (C-type lectin domain family member 9A) is also known as DNGR1 (Dendritic cell natural killer lectin group receptor 1). It is a type II membrane protein with a single extracellular C-type lectin domain. Clec9A is a dendritic cell subtype-restricted C-type lectin-like receptor. Clec9A is selectively expressed on plasmacytoid dendritic cells and CD8<sup>+</sup> myeloid dendritic cells. Clec9A reportedly serves as a receptor for necrotic cells. It can mediate the cross-presentation of dead-cell associated antigens in a Syk-dependent manner.



**Multicolor flow cytometric analysis of mouse CD370 (Clec9A) expression in spleen cells.** BALB/c splenocytes were simultaneously stained with Alexa Fluor® 700 Rat Anti-Mouse CD4 (Cat. No. 557956), APC Rat Anti-Mouse CD8 (Cat. No. 553035), FITC Hamster Anti-Mouse CD11c (Cat. No. 553801) antibodies and either PE Rat IgG2a κ Isotype Control (Cat. No. 553930; Left Plot) or PE Rat Anti-Mouse CD370 (Clec9A) antibody (Cat. No. 562734; Right Plot). Two color flow cytometric dot blots showing the correlated expression pattern of CD11c and Clec9A (or Ig Isotype control staining) were generated from CD8<sup>+</sup> CD4<sup>-</sup> gated events with the forward and side light-scatter characteristics of viable cells. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

**Preparation and Storage**

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

**Application Notes****Application**

Flow cytometry	Routinely Tested
----------------	------------------

**BD Biosciences**

bdbiosciences.com

United States 877.232.8995 Canada 866.979.9408 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

*Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.*

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

562734 Rev. 2



## Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
553930	PE Rat IgG2a, $\kappa$ Isotype Control	0.1 mg	R35-95
554656	Stain Buffer (FBS)	500 mL	(none)
557956	Alexa Fluor® 700 Rat Anti-Mouse CD4	0.1 mg	RM4-5
553035	APC Rat Anti-Mouse CD8a	0.1 mg	53-6.7
553801	FITC Hamster Anti-Mouse CD11c	0.5 mg	HL3
554657	Stain Buffer (BSA)	500 mL	(none)

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
5. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
6. Please refer to [www.bdbiosciences.com/us/s/resources](http://www.bdbiosciences.com/us/s/resources) for technical protocols.

### References

Caminschi L, Proietto AL, Ahmet F, et al. The dendritic cell subtype restricted C-type lectin Clec9A is a target for vaccine enhancement. *Blood*. 2008; 112(8):3264-3273. (Immunogen)

Huysamen C, Willment JA, Dennehy KM, Brown GD. CLEC9A is a novel activation C-type lectin-like receptor expressed on BDCA3+ dendritic cells and a subset of monocytes. *J Biol Chem*. 2008; 283(24):16693-16701. (Biology)

Sancho D, Joffre OP, Keller AM, et al. Identification of a dendritic cell receptor that couples sensing of necrosis to immunity. *Nature*. 2009; 458(7240):899-903. (Biology)