

Technical Data Sheet

Oligo Mouse Anti-Human CD44

Product Information

Material Number:	940251
Size:	25 Tests
Clone:	L178
Alternative Name:	Pgp-1; H-CAM; Hermes; ECMR-III; HUTCH-1; gp80-95; Ly-24; CSPG8; HCELL
Reactivity:	Human (Tested in Development)
Isotype:	Mouse BALB/c IgG1, κ
Immunogen:	Human TCR $\gamma\delta^+$ Thymocytes
Application:	Single Cell 3' Sequencing (Qualified)
Barcode Sequence:	GTGATTGATTAGGACAGTTCGTTGCTTAGTAGTGGG
SeqID:	AHS0167
Volume Per Test:	2 μ l
Entrez Gene ID:	960
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.
Regulatory Status:	RUO

Description

The L178 monoclonal antibody specifically binds to CD44 which is also known as Homing-associated cell adhesion molecule (H-CAM) or Phagocytic glycoprotein 1 (Pgp-1). CD44 is a single-pass type I transmembrane glycoprotein that exists in a variety of isoforms also known as variant CD44 (CD44v) or CD44R forms. CD44 is synthesized as a 37 kDa polypeptide that is converted to an 80-95 kDa form by glycosylation via N- and O-linkages, or to a 180-200 kDa form by the addition of chondroitin sulfate. It is expressed on leucocytes, erythrocytes, epithelial cells and weakly on platelets. Depending on the cell type, CD44 may participate in a variety of functions including cell adhesion, motility and cell activation.

Application Notes

The antibody was conjugated to an oligonucleotide that contains an antibody clone-specific barcode (ABC) flanked by a poly-A tail on the 3' end and a PCR handle (PCR primer binding site) on the 5' end. The ABC for this antibody was designed to be used with other BD AbSeq oligonucleotides conjugated to other antibodies. All AbSeq ABC sequences were selected in silico to be unique from human and mouse genomes, have low predicted secondary structure, and have high Hamming distance within the BD AbSeq portfolio, to allow for sequencing error correction and unique mapping. The poly-A tail of the oligonucleotide allows the ABC to be captured by the BD Rhapsody™ system. The 5' PCR handle allows for efficient sequencing library generation for Illumina sequencing platforms.

NOTE: The BD Rhapsody Single-Cell Analysis System must be used with the BD Rhapsody Express Instrument.

Preparation and Storage Section

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography and conjugated to BD AbSeq oligonucleotide under optimal conditions.

Recommended Assay Procedure

Put all BD AbSeq Reagents to be pooled into a Latch Rack for 500 μ l Tubes (Thermo Fisher Scientific Cat. No. 4900). Arrange the tubes so that they can be easily uncapped and re-capped with an 8-Channel Screw Cap Tube Capper (Thermo Fisher Scientific Cat. No. 4105MAT) and the reagents aliquoted with a multi-channel pipette.

BD AbSeq tubes should be centrifuged for ≥ 30 seconds at 400 \times g to ensure removal of any content in the cap/tube threads prior to the first opening.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS) RUO	500 mL	
633701	Single-Cell Analysis System RUO	1 Each	
564219	Human BD Fc Block™ RUO	50 mg	
564220	Human BD Fc Block™ RUO	0.25 mg	

Product Notices

1. This reagent has been pre-diluted for use at the recommended volume per test. Typical use is 2 µl for 1 × 10⁶ cells in a 200-µl staining reaction.
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.
5. Illumina is a trademark of Illumina, Inc.
6. This product is covered by one or more of the following patents: US 8,835,358; US 9,290,808; US 9,290,809; US 9,315,857; US 9,567,645; US 9,567,646; US 9,598,736; US 9,708,659; and US 9,816,137. This product, and only in the amount purchased by buyer, may be used solely for buyer's own internal research, in a manner consistent with the accompanying product literature. No other right to use, sell or otherwise transfer (a) this product, or (b) its components is hereby granted expressly, by implication or by estoppel. Diagnostic uses require a separate license.
7. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
8. Please refer to bd.com/genomics-resources for technical protocols.

References

Brown TA, Bouchard T, St John T, Wayner E, Carter WG. Human keratinocytes express a new CD44 core protein (CD44E) as a heparan-sulfate intrinsic membrane proteoglycan with additional exons. *J Cell Biol.* 1991 April; 113(1):207-221.

Carter WG, Wayner EA. Characterization of the class III collagen receptor, a phosphorylated, transmembrane glycoprotein expressed in nucleated human cells. *J Biol Chem.* 1998 March; 263(9):4193-4201.

Denning SM, Le PT, Singer KH, Haynes BF. Antibodies against the CD44 p80, lymphocyte homing receptor molecule augment human peripheral blood T cell activation.. *J Immunol.* 1990 January; 144(1):7-15.

Denning SM, Telen MJ, Hale LP, Liao HX, Haynes BF. CD44 and CD44R Cluster Report . In: :1713-1719.

Gallatin WM, Wayner EA, Hoffman PA, St John T, Butcher EC, Carter WG. Structural homology between lymphocyte receptors for high endothelium and class III extracellular matrix receptor. *Proc Natl Acad Sci U S A.* 1989 June; 86(12):4654-4658.

Jalkanen S, Jalkanen M, Bargatze R, Tammi M, Butcher EC. Biochemical properties of glycoproteins involved in lymphocyte recognition of high endothelial venules in man. *J Immunol.* 1988 September; 141(5):1615-1623.

Nagler A, Lanier LL, Cwirla S, Phillips JH. Comparative studies of human FcRIII-positive and negative natural killer cells.. *J Immunol.* 1989; 143(10):3183-91.

Shimizu Y, Van Seventer GA, Siraganian R, Wahl L, Shaw S. Dual role of the CD44 molecule in T cell adhesion and activation. *J Immunol.* 1989 October; 143(8):2457-2463.

St John T, Meyer J, Idzerda R, Gallatin WM. Expression of CD44 confers a new adhesive phenotype on transfected cells. *Cell.* 1990 January; 60(1):45-52.

BD Biosciences

bdbiosciences.com

United States 877.232.8995	Canada 888.268.5430	Europe 32.53.720.550	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 0800.771.7157
-------------------------------	------------------------	-------------------------	-----------------------	------------------------------	--

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

©2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.

