

Technical Data Sheet

Oligo Mouse Anti-Human CD72

Product Information

Material Number:	940293
Size:	25 Tests
Clone:	J4-117
Alternative Name:	Lyb-2; LYB2
Reactivity:	Human (Tested in Development)
Isotype:	Mouse IgG2b, κ
Application:	Single Cell 3' Sequencing (Qualified)
Barcode Sequence:	TAGAGTCGTTATCGTTGAGCGTATGAGCGTAGGCCT
SeqID:	AHS0215
Volume Per Test:	2 µl
Workshop No.:	VI CD72.2
Entrez Gene ID:	971
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.
Regulatory Status:	RUO

Description

The J4-117 monoclonal antibody specifically recognizes CD72. CD72 is a 39-43 kDa type II integral membrane protein that belongs to the C-type lectin superfamily. CD72 is expressed on B cells from early precursor stages through mature B cells. CD72 is not expressed on plasma cells. CD72 is also expressed on follicular dendritic reticulum cells and some macrophages. CD72 has been reported to bind to CD5, a glycoprotein expressed on all mature T cells and a subpopulation of B cells, suggesting a role in regulating T- and B-cell activation and proliferation.

Application Notes

The antibody was conjugated to an oligonucleotide that contains an antibody clone-specific barcode (ABC) flanked by a poly-A tail on the 3' end and a PCR handle (PCR primer binding site) on the 5' end. The ABC for this antibody was designed to be used with other BD AbSeq oligonucleotides conjugated to other antibodies. All AbSeq ABC sequences were selected in silico to be unique from human and mouse genomes, have low predicted secondary structure, and have high Hamming distance within the BD AbSeq portfolio, to allow for sequencing error correction and unique mapping. The poly-A tail of the oligonucleotide allows the ABC to be captured by the BD Rhapsody™ system. The 5' PCR handle allows for efficient sequencing library generation for Illumina sequencing platforms.

NOTE: The BD Rhapsody Single-Cell Analysis System must be used with the BD Rhapsody Express Instrument.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography and conjugated to BD AbSeq oligonucleotide under optimal conditions.

Recommended Assay Procedure

Put all BD AbSeq Reagents to be pooled into a Latch Rack for 500 µL Tubes (Thermo Fisher Scientific Cat. No. 4900). Arrange the tubes so that they can be easily uncapped and re-capped with an 8-Channel Screw Cap Tube Capper (Thermo Fisher Scientific Cat. No. 4105MAT) and the reagents aliquoted with a multi-channel pipette. BD AbSeq tubes should be centrifuged for ≥ 30 seconds at 400 × g to ensure removal of any content in the cap/tube threads prior to the first opening.

Suggested Companion Products

Catalog Number	Name	Size
554656	Stain Buffer (FBS)	500 mL
633701	Single-Cell Analysis System	1 Each
564219	Human BD Fc Block™	50 mg

Product Notices

1. This reagent has been pre-diluted for use at the recommended volume per test. Typical use is 2 µl for 1 × 10⁶ cells in a 200-µl staining reaction.
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.
5. Illumina is a trademark of Illumina, Inc.
6. This product is covered by one or more of the following patents: US 8,835,358; US 9,290,808; US 9,290,809; US 9,315,857; US 9,567,645; US 9,567,646; US 9,598,736; US 9,708,659; and US 9,816,137. This product, and only in the amount purchased by buyer, may be used solely for buyer's own internal research, in a manner consistent with the accompanying product literature. No other right to use, sell or otherwise transfer (a) this product, or (b) its components is hereby granted expressly, by implication or by estoppel. Diagnostic uses require a separate license.
7. Please refer to <http://regdocs.bd.com> to access safety data sheets (SDS).
8. Please refer to bd.com/genomics-resources for technical protocols.

References

Landolfi MMT, Parnes JR. CD72 Workshop Panel report. In: Kishimoto T. Tadimitsu Kishimoto .. et al., ed. Leucocyte typing VI : white cell differentiation antigens : proceedings of the sixth international workshop and conference held in Kobe, Japan, 10-14 November 1996. New York: Garland Pub.; 1997; :162-164.

Pesando JM, Stucki MA. AFTR: a fifth human B-cell-specific surface antigen. Hum Immunol. 1990 March; 37(3):193-207.

Schlossman SF, Stuart F. Schlossman .. et al., ed. Leucocyte typing V : white cell differentiation antigens : proceedings of the fifth international workshop and conference held in Boston, USA, 3-7 November, 1993. Oxford: Oxford University Press; 1995; .

BD Biosciences

bdbiosciences.com

United States
877.232.8995

Canada
888.268.5430

Europe
32.53.720.550

Japan
0120.8555.90

Asia Pacific
65.6861.0633

Latin America/Caribbean
0800.771.7157



For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

©2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.