

Technical Data Sheet

BV711 Mouse Anti-Human EphA2

Product Information

Material Number:	748149
Size:	50 µg
Clone:	371805
Alternative Name:	EPHA2; ECK; ARCC2; CTPA; CTPP1; CTRCT6
Reactivity:	Human (Tested in Development)
Isotype:	Mouse IgG2a, κ
Immunogen:	Human EphA2 Recombinant Protein
Application:	Flow cytometry (Qualified)
Concentration:	0.2 mg/ml
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.
Regulatory Status:	RUO

Description

The 371805 monoclonal antibody specifically recognizes Ephrin type-A receptor 2 (EphA2). EphA2 is a ~130 kDa type I transmembrane glycoprotein that is encoded by EPHA2 (EPH receptor A2) which belongs to ephrin receptor subfamily within the receptor tyrosine kinase (RTK) family. EphA2 is also known as Epithelial cell kinase (ECK) or Epithelial cell receptor protein tyrosine kinase. The extracellular region of EphA2 contains an N-terminal globular domain, a cysteine-rich domain and 2 fibronectin type III repeats, whereas the cytoplasmic domain contains two tyrosine residues adjacent to the membrane that function as major autophosphorylation sites and a kinase domain. EphA2 is variably expressed by epithelial cells, dendritic cells, Langerhans cells, keratinocytes, and endothelial cells. EphA2 functions as a receptor for glycosphosphatidylinositol (GPI) membrane-linked members of the Ephrin-A family, including Ephrins A1-A5. EphA2 is involved in regulating cellular growth, adhesion, migration, survival, and plays a role in angiogenesis. Its expression may be upregulated on vascular endothelium in certain breast, prostate, and colon cancers as well as on some metastatic tumor cells.

The antibody was conjugated to BD Horizon™ BV711 which is part of the BD Horizon Brilliant™ Violet family of dyes. This dye is a tandem fluorochrome of BD Horizon BV421 with an Ex Max of 405-nm and an acceptor dye with an Em Max at 711-nm. BD Horizon BV711 can be excited by the violet laser and detected in a filter used to detect Cy™5.5 / Alexa Fluor® 700-like dyes (eg, 712/20-nm filter). Due to the excitation and emission characteristics of the acceptor dye, there may be moderate spillover into the Alexa Fluor® 700 and PerCP-Cy5.5 detectors. However, the spillover can be corrected through compensation as with any other dye combination.

Preparation and Storage Section

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with BD Horizon BV711 under optimal conditions that minimize unconjugated dye and antibody.

Recommended Assay Procedure

For optimal and reproducible results, BD Horizon Brilliant Stain Buffer should be used anytime two or more BD Horizon Brilliant dyes (including BD OptiBuild Brilliant reagents) are used in the same experiment. Fluorescent dye interactions may cause staining artifacts which may affect data interpretation. The BD Horizon Brilliant Stain Buffer was designed to minimize these interactions. More information can be found in the Technical Data Sheet of the BD Horizon Brilliant Stain Buffer (Cat. No. 563794).

Suggested Companion Products

Catalog Number	Name	Size	Clone
349202	Lysing Solution 10X Concentrate IVD	100 NA	
564219	Human BD Fc Block™ RUO	50 mg	
554656	Stain Buffer (FBS) RUO	500 mL	
554657	Stain Buffer (BSA) RUO	500 mL	
563794	Brilliant Stain Buffer RUO	100 Tests	
555899	Lysing Buffer RUO	100 mL	

Product Notices

1. This antibody was developed for use in flow cytometry.
2. The production process underwent stringent testing and validation to assure that it generates a high-quality conjugate with consistent performance and specific binding activity. However, verification testing has not been performed on all conjugate lots.
3. Researchers should determine the optimal concentration of this reagent for their individual applications.
4. An isotype control should be used at the same concentration as the antibody of interest.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at wwwbdbiosciences.com/colors.
7. Please refer to wwwbdbiosciences.com/us/s/resources for technical protocols.
8. BD Horizon Brilliant Stain Buffer is covered by one or more of the following US patents: 8,110,673; 8,158,444; 8,575,303; 8,354,239.
9. BD Horizon Brilliant Violet 711 is covered by one or more of the following US patents: 8,110,673; 8,158,444; 8,227,187; 8,455,613; 8,575,303; 8,354,239.
10. Cy is a trademark of GE Healthcare.
11. Alexa Fluor® is a registered trademark of Life Technologies Corporation.

References

- Brantley DM, Cheng N, Thompson EJ, et al. Soluble Eph A receptors inhibit tumor angiogenesis and progression in vivo. *Oncogene*. 2002; 21(46):7011.
- Cheng N, Brantley DM, Liu H, et al. Blockade of EphA receptor tyrosine kinase activation inhibits vascular endothelial cell growth factor-induced angiogenesis. *Mol Cancer Res*. 2002; 1(1):2-11.
- de Saint-Vis B, Bouchet C, Gautier G, Valladeau J, Caux C, Garrone P. Human dendritic cells express neuronal Eph receptor tyrosine kinases: role of EphA2 in regulating adhesion to fibronectin. *Blood*. 2003; 102(13):4431-4440.
- Liao-Chan S, Daine-Matsuoka B, Heald N, et al. Quantitative assessment of antibody internalization with novel monoclonal antibodies against Alexa fluorophores. *PLoS ONE*. 2015; 10(4):e0124708.
- Lindberg RA, Hunter T. cDNA cloning and characterization of eck, an epithelial cell receptor protein-tyrosine kinase in the eph/elk family of protein kinases. *Mol Cell Biol*. 1990; 10(12):6316-24.
- Udayakumar D, Zhang G, Ji Z, Njauw CN, Mroz P, Tsao H. EphA2 is a critical oncogene in melanoma. *Oncogene*. 2011; 30(50):4921-4929.
- Udyavar AR, Wooten DJ, Hoeksema M, et al. Novel Hybrid Phenotype Revealed in Small Cell Lung Cancer by a Transcription Factor Network Model That Can Explain Tumor Heterogeneity. *Cancer Res*. 2017; 77(5):1063-1074.

BD Biosciences

bdbiosciences.com

United States
877.232.8995

Canada
888.268.5430

Europe
32.53.720.550

Japan
0120.8555.90

Asia Pacific
65.6861.0633

Latin America/Caribbean
0800.771.7157

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for a patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

©2020 BD. All rights reserved. Unless otherwise noted, BD, the BD Logo and all other trademarks are the property of Becton, Dickinson and Company or its affiliates.

